

CLEAN SYNTHETIC VECTORS, PLASMIDS, TRANSGENIC PLANTS  
AND PLANT PARTS CONTAINING THEM, AND METHODS FOR OBTAINING  
THEM.

5 This application claims the priority of PCT  
Application No. PCT/IB00/01243, filed September 4, 2000, and  
published in English on March 15, 2001, which application  
claims the priority of French Patent Application No. 99111112,  
10 filed September 3, 1999.

The present invention relates to clean synthetic  
vectors, intended notably for use for genetic transformation  
in the field of plant biotechnology.

15 In general, the vectors are known in the field of  
biotechnology and genetic manipulation. The vectors which are  
currently used most commonly for genetic transformation, and  
in particular in the field of plant biotechnology, present  
several disadvantages in their use, however. In actual fact,  
20 and notably for plant transgenesis, use has often been made of  
a vector called pBin19 (Frisch et al., 1995). The nucleotide  
sequence of this binary plasmid pBin19 is entirely known. The  
problem, however, is that this plasmid is of large size (11.8  
kbp) and that it contains useless elements (more than half of  
25 pBin19) which are intolerable from a regulatory point of view,  
or even detrimental to good replication. Moreover, the  
selection cassette of this plasmid is located near the right  
border of the T-DNA. Thus, when the T-DNA is broken after the  
selection cassette, it will not be possible to make any  
30 selection in order to retain solely the plants possessing the  
expression cassette of the desired gene.

The expressions used in the description and the  
claims have the following meaning :

- "vector" means an expression system, for example  
35 DNA-coated projectiles, nucleic-acid-based transit vehicles,  
nucleic acid molecules adapted to deliver nucleic acid, and

autonomous self-replicating circular DNA, for example plasmids, cosmids, phagemids, etc. If a micro-organism or a recombinant cell culture is described as host of an "expression vector", this can also include extrachromosomal circular DNA (such as for example mitochondrial or chloroplast DNA), DNA which has been integrated into the host chromosome(s), where the vector can be either replicated in a stable manner by the cells during mitosis as an autonomous structure, integrated into the genome of the host, or maintained in the nucleus or cytoplasm of the host.

- usually the vectors used for the genetic transformation exist in the form of plasmids. In this case, the "plasmid" is a molecule of autonomous circular DNA capable of replication in a cell. If a micro-organism or recombinant cell culture is described as the host of an "expression" plasmid, this comprises both extrachromosomal circular DNA molecules and DNA which has been integrated into the host chromosome(s). If the plasmid is maintained by a host cell, the plasmid is either replicated in a stable manner by the cells during mitosis as an autonomous structure, or integrated into the genome of the host;

- "clean" means that the vector comprises only sequences that are indispensable for its functionality and carries a nucleic acid sequence which comprises only elements that are indispensable for the expression of the host cell;

- "nucleic acid" means DNA or RNA ;

- "nucleic acid sequence" means a single- or double-stranded oligomer or polymer of nucleotide bases read from the 5' end towards the 3' end, and comprises self-replicating plasmids, genes, DNA or RNA polymers, which may or may not be infectious, and DNA, or RNA, either functional or non-functional. In the nucleotide notation used in the present application, unless specifically stated, the left end of a single-stranded nucleotide sequence is the 5' end;

- "derived nucleic acid sequence" means that the sequence derives directly or indirectly from the sequence

referred to, for example by substitution, deletion, addition, mutation, fragmentation, and/or synthesis of one or more nucleotides;

5 - "promoter" means a nucleic acid region which is upstream of the translation initiation codon and which is involved in the recognition and binding of RNA polymerase and other transcription proteins;

- "plant promoter" is a promoter capable of initiating transcription in plant cells;

10 - "constitutive promoter" is a promoter capable of expressing nucleic acid sequences operationally bound to said promoter, in all or practically all the tissues of the host organism, during the whole development of said organism ;

15 - "tissue specific promoter" is a promoter capable of selectively expressing nucleic acid sequences operationally bound to said promoter, in certain specific tissues of the host organism;

20 - "operationally bound" means the binding of a functional or regulatory element, for example a promoter, to the nucleic acid sequence, or gene, to be expressed which codes for a protein to be produced, in such a way that this element influences the transcription of the bound nucleic acid sequence;

25 - "expression cassette" means nucleotide sequences capable of directing the expression of a nucleic acid sequence, or of a gene, coding for a polypeptide to be produced in a host organism compatible with such sequences. Such cassettes include at least one promoter and a transcription termination signal, and optionally other factors  
30 necessary or useful for the expression ;

- "heterologous sequence" or "heterologous nucleic acid sequence" means a sequence originating from a source, or from a species, foreign to the environment thereof, or if it originates from the same environment, which has been modified  
35 relative to its original form. The modification of the nucleic acid sequence can take place for example by treatment of the

nucleic acid with a restriction enzyme in order to generate a nucleic acid fragment which can be operationally bound to a promoter. The modification can also take place by means of techniques such as directed mutagenesis;

5       - "box" means a nucleic acid sequence to which a regulatory function is attributed;

      - "like" means that the box, and/or the nucleic acid sequence with which this term is associated, involves a certain sequence identity or consensus with a box and/or a known nucleic acid sequence, called a reference sequence, preferably a sequence identity of at least 50%, more preferably a sequence identity of at least 75%, and more particularly a sequence identity of at least 90% with the reference sequence. The percentage of sequence identity is calculated on the basis of a comparison window of at least 6 nucleotide bases. The determination of a comparison window can be carried out using sequence alignment algorithms to determine a homology with a reference sequence, for example the local homology algorithm, the homology alignment algorithm, and the similarity search algorithm, these algorithms existing also in computerized form, known by the names GAP, BESTFIT, FASTA and TFASTA. The percentage of sequence identity is obtained by comparing the reference sequence with the box and/or the nucleic acid sequence;

25       - "situated" means the position on a nucleic acid sequence of an identified element, such as a "box", a restriction site, or a codon having a particular function. The position which is given by a figure refers to the position of the start of the element in the nucleic acid sequence, in the direction of reading of the latter, i.e. in the direction 5'→3';

30       - "transgenic plant" means a plant which has been obtained by genetic manipulation techniques, and covers the whole plants obtained, their progeny, and the plant organs, for example the roots, stems and leaves, obtained by these techniques. The transgenic plants according to the present

invention can have different levels of ploidy, and can notably be polyploid, diploid, and haploid;

- "propagule" means a cluster or association of plant cells, which may or may not be structured, allowing the  
5 regeneration of a whole plant, for example explants, calli, stems, leaves, roots, cuttings, and even seeds.

The applicant of the present invention has succeeded, surprisingly, in producing clean synthetic vectors, in particular binary plasmids, of completely known nucleotide  
10 sequence, of small size, allowing the aforesaid disadvantages to be alleviated, and notably presenting a high replication rate relative to the existing vectors most commonly used. Furthermore, the applicant has succeeded at the same time in  
15 producing a range of vectors in such a way as to be able to choose the one which it is convenient to use according to the application envisaged and the environment of its use, and thus in such a way as to be able to better control the rate of expression of a gene to be expressed, coding for a polypeptide to be produced.

20 In addition, in some of the vectors according to the invention, each of the functional elements or components can be isolated by simple enzymatic digestion.

An object of the present invention is therefore a clean synthetic vector containing only the elements  
25 indispensable to its functionality and to the transgenesis of a cell, and notably of a plant cell.

According to a preferred embodiment, the vector comprises, as elements which are indispensable to its functionality and to the transgenesis of a cell, and which are  
30 operationally bound:

- at least one nucleic acid sequence coding for at least one first origin of replication, preferably an ori RK2, and more preferably at least one ori V of pRK2 of Escherichia coli with a broad host range;

35 - at least one nucleic acid sequence coding for a selection agent, preferably an antibiotic resistance gene,

more preferably the npt III gene conferring resistance to kanamycin in bacteria;

- a trfA locus coding for at least one protein allowing an increase in the replication rate of the plasmid, preferably originating from pRK2, and more preferably coding for the proteins P285 and P382.

According to a more preferred embodiment the vector comprises the nucleic acid sequence identified by the number SEQ.ID01. Still more preferably, the vector consists of a single plasmid pMRT1105 whose nucleic acid sequence is identified by the number SEQ.ID01.

According to another preferred embodiment, the vector includes at least one nucleic acid sequence coding for a second origin of replication, preferably an ori of Escherichia coli, and more preferably an ori ColEI. More preferably, the vector comprises the nucleic acid sequence identified by the number SEQ.ID02, and still more preferably, the vector consists of a single plasmid pMRT1106 whose nucleic acid sequence is identified by the number SEQ.ID02.

Preferably, the synthetic vector according to the invention comprises a region (200 bp or less) made up of at least one nucleic acid sequence containing several unique enzymatic restriction sites called collectively «multiple cloning sites» (MCS).

According to a preferred embodiment, the vector also includes a nucleic acid sequence coding for a T-DNA comprising a right border RB and a left border LB, allowing the vector to act as a binary plasmid.

Preferably, the MCS is situated near (within 300 bp of) the right border, RB, of the T-DNA prior to the cloning of an insert into the MCS.

According to another preferred embodiment, the vector also includes a nucleic acid sequence coding for at least one expression promoter and at least one transcription terminator situated between the left border LB and the right border RB of the T-DNA. More preferably, the expression promoter is chosen

from the group consisting of the constitutive promoters, the inducible promoters, the specific promoters, and preferably chosen from the plant expression promoters. Still more preferably, the expression promoter is chosen from the group consisting of the 35S CaMV promoter, the ep35S of CaMV, the pea plastocyanin gene promoter, its "enhancer" zones and derivatives, the wheat "High Molecular Weight Glutenin" (HMWG) promoter, the "Cassava Vein Mosaic Virus" CsVMV promoter, the "Commelina Yellow Mottle Virus" CoYMV promoter, the chimeric promoters of the CsVMV and CoYMV promoters, and their derivatives.

Preferably, the expression terminator is chosen from the functional terminators in a plant cell, and is preferably a 35S or nos terminator.

According to yet another preferred embodiment, the vector comprises at least one nucleic acid sequence coding for a selection agent that is functional in a plant cell, preferably at least one acid sequence coding for an antibiotic resistance gene, and/or a herbicide resistance gene.

Preferably, the sequence coding for a selection agent is a sequence coding for the bar («bialaphos resistance») resistance gene or pat («phosphinothricin acetyltransferase») gene, or else a sequence coding for the mutant or wild-type resistance gene nptII. More preferably still, the nucleic acid sequence coding for the selection agent is situated near the left border of the T-DNA.

According to another preferred embodiment of the present invention, the vector includes at least one expression cassette comprising an expression-promoting nucleic acid sequence operationally bound to a nucleic acid sequence to be expressed, coding for a polypeptide to be produced, itself bound to a transcription termination nucleic acid sequence. Preferably, the polypeptide to be produced is an enzyme or protein or derivative of the latter having activity in vitro and/or in man and/or in animals, said activity comprising digestive, pancreatic, biliary, antiviral, anti-inflammatory,

pulmonary, anti-microbial, nutritional, cosmetic, structural, blood, cardiovascular, ophthalmic, antigenic, immunostimulatory and cerebral activity. Examples of such proteins are for example the insulins, interferons, gastric, pancreatic or biliary lipases, elastases, antiproteases such as alpha-1 antitrypsin, structural proteins such as collagen, transferrins such as lactoferrin, proteins derived from blood, such as haemoglobin, human albumin and the blood cofactors, and antioxidants such as superoxide dismutase,

10 According to a particularly preferred embodiment of the invention, the vector is presented in the form of a binary, linear or circular plasmid, chosen from the group consisting of the nucleic acid sequences identified by the numbers SEQ.ID03, SEQ.ID04, SEQ.ID05, SEQ.ID06, SEQ.ID07, 15 SEQ.ID08, SEQ.ID09, SEQ.ID10, SEQ.ID11, SEQ.ID12, SEQ.ID13, SEQ.ID14, SEQ.ID15, SEQ.ID16, SEQ.ID17, SEQ.ID18, SEQ.ID19, SEQ.ID20, SEQ.ID21 and SEQ.ID22.

20 According to a particularly advantageous embodiment, each functional component of the vector can be cleaved independently of the other components. Preferably, each functional component can be cleaved independently of the other components by enzymatic digestion at the level of a first unique restriction site and a second unique restriction site which are present in 1 vector.

25 Another object of the present invention is an isolated nucleic acid sequence, characterized in that it corresponds to a nucleic acid sequence chosen from the group consisting of the nucleic acid sequences identified by the numbers SEQ.ID01, SEQ.ID02, SEQ.ID03, SEQ.ID04, SEQ.ID05, 30 SEQ.ID06, SEQ.ID07, SEQ.ID08, SEQ.ID09, SEQ.ID10, SEQ.ID11, SEQ.ID12, SEQ.ID13, SEQ.ID14, SEQ.ID15, SEQ.ID16, SEQ.ID17, SEQ.ID18, SEQ.ID19, SEQ.ID20, SEQ.ID21 and SEQ.ID22.

35 Yet another object of the present invention is a cell containing a vector or a nucleic acid sequence such as described earlier. The cell is preferably a plant cell.



Yet another object of the present invention is a transgenic plant having stably integrated in its genome a vector or a nucleic acid sequence such as described earlier. The plant is preferably chosen from the dicotyledon species, such as potato, tobacco, cotton, lettuce, tomato, melon, cucumber, pea, rape, beetroot or sunflower, or the monocotyledon species, such as wheat, barley, oats, rice or maize.

Yet another object of the present invention is a propagule of a transgenic plant such as described earlier. The propagule is preferably a seed.

Yet another object of the present invention is a method for expression of a nucleic acid sequence, or gene, coding for a polypeptide to be produced, in a cell, characterized in that it comprises the stages consisting of:

- transforming the cell with a vector or a nucleic acid sequence such as described earlier;

- making a culture of the cell under conditions allowing the expression of a nucleic acid sequence, or gene, coding for the polypeptide to be produced. The cell is preferably a prokaryotic or eukaryotic cell. More preferably, the cell is a cell chosen from the group consisting of microbial cells, fungal cells, insect cells, animal cells and plant cells. Still more preferably, the cell is a plant cell.

Yet another object according to the present invention is a method for obtaining a transgenic plant or a propagule such as described earlier, characterized in that it comprises the stages consisting of:

- transforming a plant cell with a vector comprising or a nucleic acid sequence according to any one of Claims 1 to 24 ;

- selecting the plant cell having the vector or nucleic acid sequence integrated therein;

- propagating the transformed and selected plant cell; either by culture or by regeneration of whole chimeric or transgenic plants.

Thus, the new synthetic vectors, and preferably, binary plasmids, according to the invention have only regions indispensable for the functionality of the vector and for transgenesis. The applicant has found that they have a high replication rate. In addition, they can have a selection cassette near the left border, as well as rare restriction sites in their «polylinkers» (multiple cloning sites).

#### DESCRIPTION OF THE FIGURES

The invention will be understood better from the detailed description of the different embodiments which is given below in the form of examples, which are not limiting, and with reference to the drawing in the annex, in which:

- Figure 1 schematically represents a preferred minimal vector according to the present invention in the form of a plasmid identified by the reference pMRT1105, and with a length of 3508 base pairs;
- Figure 2 represents a schematic chart of a variant of the vector of Figure 1 in the form of a plasmid identified by the reference pMRT1106, and with a length of 4098 base pairs ;
- Figure 3 represents a schematic chart of a preferred vector according to the present invention, in the form of a binary plasmid identified by the reference pMRT1118, with a length of 5971 base pairs, and including a T-DNA comprising a selection cassette coding for resistance to an antibiotic, and a multiple cloning site (MCS);
- Figure 4 represents a preferred embodiment of the vector according to Figure 3, the multiple cloning site (MCS) including yet other unique sites, the vector being in the form of a binary plasmid identified by the reference pMRT1119, and with a length of 6016 base pairs ;

Figures 5 to 22 represent other preferred embodiments of the vector according to the invention, in the form of binary plasmids identified respectively by the references pMRT1121 (6017 base pairs), pMRT1122 (6016 base pairs), pMRT1155 (6017 base pairs), pMRT1175 (6767 base pairs),

PMRT1176 (6767 base pairs), PMRT1191 (4805 base pairs),  
 PMRT1192 (8654 base pairs), PMRT1193 (9143 base pairs),  
 PMRT1195 (6865 base pairs), PMRT1196 (8654 base pairs),  
 PMRT1201 (7943 base pairs), PMRT1202 (5614 base pairs),  
 PMRT1203 (7503 base pairs), PMRT1204 (9390 base pairs),  
 PMRT1205 (7503 base pairs), PMRT1206 (9390 base pairs),  
 PMRT1210 (10003 base pairs) and PMRT1212 (8987 base pairs)  
 ;

Figure 23 graphically represents a comparison between the  
 protein expression capacities of the sythetic vectors  
 according to the present invention in comparison to a  
 known vector system ;

Figures 24 to 28 represent further embodiments of vectors  
 according to the present invention, in the form of binary  
 plasmids identified by the references PMRT1334 (9688 bp),  
 PMRT1335 (15208 bp), PMRT1336 (9285 bp), PMRT1337 (8289  
 bp), PMRT1341 (14108 bp), PMRT1342 (15077 bp).

In the figures, the abbreviations have the following  
 meanings :

- ori RK2 : ori V of pRK2 of Escherichia coli with a broad  
 host range, unstable RK2 replicon producing a low number of  
 copies;
- ori ColEI : the origin of replication ColEI of Escherichia  
 coli;
- npt III gene: coding for neomycin phosphotransferase  
 conferring resistance to kanamycin
- npt II gene: coding for neomycin phosphotransferase  
 conferring resistance to kanamycin ;
- trfA : trfA locus (1481 bp) originating from pRK2 allowing  
 the production of 2 proteins, P285 and P382, which enhance  
 the replication of the plasmid by binding to the origin of  
 replication ;
- Tnos : nos (nopaline synthetase) terminator;
- Pnos : nos (nopaline synthetase) promoter;
- LB : left border of a T-DNA ;
- RB : right border of a T-DNA ;

- MCS : multiple cloning site, S:XXX indicating the number of the starting base, E:XXX indicating the number of the end base of the MCS, the list starting with the first restriction site at the top and ending with the last restriction site at the bottom, in the direction 5'>3' ;
- polyA 35S : transcription termination (polyadenylation) signal ;
- gus gene: gene coding for beta-glucuronidase ;
- IA : the actin intron of rice ;
- PA : actin promoter of rice ;
- Phmwg : «high molecular weight glutenin» promoter of wheat ;
- bar («bialaphos resistance») gene: gene coding for the enzyme phosphinothricin acetyltransferase conferring resistance to glufosinate ;
- ep35S : «enhanced promoter» of the 35S ribosome.

#### EXAMPLES

In the detailed description which follows, all the enzymatic digestions by restriction enzymes were performed in accordance with the recommendations of the supplier New England Biolabs. The purifications with the aid of the «QIAquick Gel Extraction» and «QIAquick PCR Purification» kits were carried out in accordance with the recommendations of the supplier QIAGEN. The «Concert Rapid PCR Purification System» kit was used in accordance with the instructions of the supplier GIBCO BRL Life Technologies. The «GeneAmp PCR System 9700» thermocycler used is marketed by Perkin Elmer Applied Biosystems.

#### Example 1

##### 1. Synthesis of pMRT1105

A clean synthetic vector according to the invention is presented preferably in the form of a minimal plasmid pMRT1105 (3508 bp) and is made up of the following elements:

- ori RK2 : ori V of pRK2 of Escherichia coli with a broad host range, unstable RK2 replicon producing a small number of copies (643 bp).

- npt III gene : confers resistance to kanamycin (bacterial selection marker, 1337 bp).

- TrfA : trfA locus (1481 bp) originating from pRK2, allowing the production of 2 proteins, P285 and P382, which enhance the replication of the plasmid by binding to the origin of replication.

The plasmid pMRT1105 results from the assembling of the fragments obtained by splicing overlap extension for «ori RK2» and a «part of npt III», of the fragment corresponding to a «part of trfA» produced by PCR («polymerase chain reaction») amplification and of «parts of trfA and npt III» isolated by enzymatic digestion.

#### 1.1. Synthesis of the fragment carrying «ori RK2 and part of npt III»

The AvrII-StuI fragment (654 bp) carrying «ori RK2» (643 bp) was amplified by PCR from 5 ng of pBin19 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' AACCTAGGAAAAGACCGAGCGCCTTTGC 3' (SEQ.ID23) containing the AvrII restriction site and 5' CGGATTAATGGTAGAAGGCCTTTTCACGGGAGGGTTCGAGAAGG 3' (SEQ.ID24) possessing the StuI restriction site, in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH 9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 55°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3 min. Forty µl of the PCR reaction medium were then subjected to the action of 12.5 U of the Klenow fragment (New England Biolabs) in the presence of 2 µl of each of the dNTPs at 10 mM. The reaction was carried out at 37°C for 10 min. The PCR product so treated was then isolated by 2% agarose gel electrophoresis in TBE buffer (90 mM Tris-HCl, 2 mM Na<sub>2</sub>-EDTA,

90 mM boric acid, pH 8.0) and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA was recovered in 30 µl of H<sub>2</sub>O.

The StuI-BstXI fragment (363 bp) carrying «part of npt III» (344 bp) was amplified and treated in the same way as the «ori RK2» fragment except that the 2 oligodeoxynucleotides used are 5' TGAAAGGCCTTCTACCATTAATCCGCGATAAACCCAGCGAACC 3' (SEQ.ID25) containing an StuI restriction site and 5' ATGCATCCAAAATTTTGGTAGAATTTACAAGCTATAAGGTTATTGTCCTGGG 3' (SEQ.ID26) possessing the BstXI restriction site.

The 2 fragments carrying «ori RK2» and «part of npt III» were then assembled by splicing overlap extension in order to obtain the fragment «ori RK2 and part of npt III». To do this, a PCR amplification was performed from 7.5 µl of each of the treated PCR products corresponding to «ori RK2» and «part of npt III» with the aid of 20 pmoles of each of the oligodeoxynucleotides 5' AACCTAGGAAAAGACCGAGCGCCTTTGC 3' (SEQ.ID23) and 5' ATGCATCCAAAATTTTGGTAGAATTTACAAGCTATAAGGTTATTGTCCTGGG 3' (SEQ.ID26) in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH 9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of ELONGASE enzyme in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 62°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3 min. Forty µl of the PCR reaction medium were then subjected to 1.5% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA, corresponding to the fragment carrying «ori RK2» and «part of npt III», was recovered in 30 µl of H<sub>2</sub>O. Next, 27 µl of this DNA were hydrolysed by BstXI followed by AvrII, then purified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of H<sub>2</sub>O.

### 1.2. Synthesis of the fragment carrying a «part of trfA»

The NdeI-AvrII fragment (295 bp) carrying «part of trfA» (295 bp) was amplified and treated in the same way as the fragment «ori RK2» except that the 2 oligodeoxynucleotides used are 5' ATCGACGAGGAAATCGTCGCTGCTTTGC 3' (SEQ.ID27) situated upstream of the NdeI site and 5' AAACCTAGGAAATGCCAGTAAAGCGCTGGC 3' (SEQ.ID28) possessing the AvrII restriction site. The DNA fragments corresponding to a «part of trfA» originating from the PCR amplification are then purified with the aid of the «QIAquick PCR Purification» kit, recovered in 30 µl of H<sub>2</sub>O, hydrolysed by AvrII and NdeI, repurified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of H<sub>2</sub>O.

### 1.3. Synthesis of the fragment carrying the «parts of trfA and npt III»

The BstXI-NdeI fragment (2196 bp) carrying the «parts of trfA and npt III» (2196 bp) was isolated from 9 µg of pBin19 DNA by enzymatic digestion by BstXI, purified with the aid of the «QIAquick PCR Purification» kit, then hydrolysed by NdeI. The DNA fragment was then isolated by 0.9% agarose gel electrophoresis in TBE buffer, purified with the aid of the «QIAquick Gel Extraction» kit and recovered in 30 µl of H<sub>2</sub>O.

### 1.4. Obtaining pMRT1105

The 3 fragments, «ori RK2 and part of npt III», «parts of trfA and npt III» and «part of trfA», were ligated. To do this, the reaction mixture containing 7.5 µl of the digested fragments corresponding to «ori RK2 and part of npt III» and «parts of trfA and npt III», and 11 µl of the digested fragment corresponding to «part of trfA», was concentrated with the SpeedVac AES1000 (SAVANT) in order to achieve a volume of 8 µl. After that, 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs) were added. The ligation was carried out by a PCR reaction made up of 200 cycles each consisting of 2 stages, one at 30°C for 30 sec. and the other

at 10°C for 30 sec., in the «GeneAmp PCR System 9700» thermocycler. The bacteria, *Escherichia coli* DH5α, which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on  
5 Luria-Bertani medium (LB, bactotryptone 10 g/l, yeast extract 5 g/l, NaCl 10 g/l, Agar 15 g/l, pH 7.5) supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The resulting plasmid selected  
10 was called pMRT1105 (3508 bp) and is represented in Figure 1. Its complete sequence SEQ.ID01 is given in the sequence listing.

## 2 Synthesis of pMRT1106

15 The plasmid pMRT1106 (4098 bp) differs from pMRT1105 by the addition of the origin of replication ColE1 of *Escherichia coli* («ori ColE1»).

The fragment, carrying «ori ColE1» (590 bp), was isolated from the plasmid pBR322 marketed by New England  
20 Biolabs. The plasmid (5 µg) was digested by NdeI, purified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of TE buffer (10 mM Tris-HCl, 1 mM Na<sub>2</sub>-EDTA, pH 8.0). The plasmid thus linearized was subjected to the action of 20 units of the Klenow fragment (New England  
25 Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl<sub>2</sub>, 6 µl of 1M dithiothreitol, 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min. The linearized plasmid so treated was purified with the aid of the «QIAquick PCR Purification» kit and recovered in 30 µl of  
30 H<sub>2</sub>O. It was then digested by AlwNI in order to isolate the fragment carrying «ori ColE1», purified with the aid of the «QIAquick PCR Purification» kit and subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction medium of 120 µl in the presence of 12 µl of T4 DNA  
35 polymerase x 10 buffer, 4 µl of 10 mM dNTP and 6 µl of BSA 1 mg/ml. The reaction was carried out at 37°C for 30 min. The



DNA fragment was then isolated by 1.2% agarose gel electrophoresis in TBE buffer, purified with the aid of the «QIAquick Gel Extraction» kit and recovered in 50  $\mu$ l of H<sub>2</sub>O. This DNA fragment was then inserted into the plasmid pMRT1105 (2  $\mu$ g) digested by StuI, purified with the aid of the «QIAquick PCR Purification» kit, dephosphorylated by 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the «QIAquick PCR Purification» kit. The ligation by PCR reaction was carried out with 10 ng of digested dephosphorylated pMRT1105 plasmid and 100 ng of DNA fragments carrying «ori ColE1» in a reaction medium of 10  $\mu$ l in the presence of 1  $\mu$ l of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec. in the «GeneAmp PCR System 9700» thermocycler. The bacteria, Escherichia coli DH5 $\alpha$ , which had previously been made competent, were transformed (Hanahan, 1983). The validity of the clones obtained was verified by a PCR test on bacterial colonies using 10 pmol of each of the 2 oligodeoxynucleotides, 5' AACCTAGGAAAAGACCGAGCGCCTTTGC 3' (SEQ.ID23) and 5' ATGCATCCAAAATTTTGGTAGAATTTACAAGCTATAAGGTTATTGTCCTGGG 3' (SEQ.ID26), in a reaction medium of 24  $\mu$ l in the presence of 22  $\mu$ l of SuperMix PCR (GIBCO BRL Life Technologies). The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 5 min., the DNA was subjected to 30 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 60°C for 30 sec. and elongation at 72°C for 30 sec. Then, in the last cycle, the elongation was continued at 72°C for 7 min. The PCR reaction medium was then subjected to 1% agarose gel electrophoresis in TBE buffer. The insertion of the sequence «ori ColE1» was visualized by the presence of a fragment of about 1.6 kbp. The plasmid DNA of the clones

obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The resultant plasmid selected was called  
5 pMRT1106 (4098 bp) and is represented in Figure 2. Its complete sequence SEQ.ID02 is given in the sequence listing.

### 3. Synthesis of pMRT1118

The plasmid pMRT1118 (5971 bp) differs from pMRT1106  
10 by the introduction of a clean transfer DNA (T-DNA) in pMRT1106. This clean T-DNA is made up of the expression cassette of the mutant nptII gene (Frisch et al., 1995) under the control of the promoter and terminator of the nopaline synthase gene (nos, Depicker et al., 1982) of *Agrobacterium*  
15 *tumefaciens* placed between the right (RB) and left border (LB) of the plasmid pTiT37 of *Agrobacterium tumefaciens* nopaline strain.

#### 3.1 Synthesis of the fragment carrying «end nptII - Tnos - LB»

##### 20 3.1.1. Obtaining the fragment carrying LB

The AvrII-SmaI fragment (173 bp) carrying LB (151 bp) was amplified by PCR from 5 ng of pBin19 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29)  
25 containing the AvrII and AatII restriction sites, and 5' CCTATGGATATCCCCCGGGGATAGCCCCAGTACATTAAAAACGTCC 3' (SEQ.ID30) possessing the SmaI restriction site, in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH 9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of eLONGASE enzyme (GIBCO BRL  
30 Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at  
35 55°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3

min. Forty  $\mu$ l of the PCR reaction medium were then subjected to the action of 12.5 U of the Klenow fragment (New England Biolabs) in the presence of 2  $\mu$ l of each of the dNTPs at 10 mM. The reaction was carried out at 37°C for 10 min. The PCR product so treated was then isolated by 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA was recovered in 50  $\mu$ l of H<sub>2</sub>O.

3.1.2. Obtaining the fragment carrying the nos terminator (Tnos)

The SmaI-BspEI fragment (288 bp) carrying the nos terminator (256 bp) was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are  
5'CTATCCCCCGGGGATATCCATAGGCCCGATCTAGTAACATAGATGAC 3'  
(SEQ.ID31) containing a SmaI restriction site and 5' GCGCACTTGGGCCCATAGCTCGACGAACGATCGTTCAAACATTTGGC 3' (SEQ.ID32), possessing the Bsp120I restriction site.

3.1.3. Obtaining the fragment carrying «end of the nptII» gene («end nptII»)

The fragment (221 bp) carrying «end nptII» (221 bp) was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are  
5' TTCGTCGAGCTATGGGCCCAAGTGCATCCCGTGGGCGAAGAACTC 3'  
(SEQ.ID33) containing a Bsp120I restriction site and 5' TTCTTGACGAGTTCTTCTGAGCGGG 3' (SEQ.ID34) downstream of BstBI.

3.1.4. Obtaining the fragment «end nptII - Tnos - LB»

The fragments, «LB», «Tnos» and «end nptII», were assembled by splicing overlap extension in 2 stages :

- assembly of the fragments «LB» and Tnos» resulting in a fragment «Tnos - LB»,
- assembly of the fragment «Tnos - LB» and of the fragment «end nptII» resulting in a fragment «end nptII - Tnos - LB».

To do this, a first PCR amplification was carried out from 10  $\mu$ l of each of the PCR products treated corresponding

to «LB» and «Tnos», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5'  
TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5'  
GCGCACTTGGGCCCATAGCTCGACGAACGATCGTTCAAACATTGGC 3' (SEQ.ID32), in  
5 the presence of 200  $\mu$ M of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH  
9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of ELONGASE enzyme  
(GIBCO BRL Life Technologies) in a final reaction medium of  
50  $\mu$ l. The PCR amplification reaction was carried out in the  
«GeneAmp PCR System 9700» thermocycler. After denaturation at  
10 94°C for 3 min., the DNA was subjected to 15 cycles each made  
up of the stages of denaturation at 94°C for 45 sec.,  
hybridization at 62°C for 45 sec. and elongation at 68°C for 1  
min. Then, in the last cycle, the elongation was continued at  
68°C for 3 min. Forty  $\mu$ l of the PCR reaction medium were then  
15 subjected to the action of 12.5 U of the Klenow fragment (New  
England Biolabs) in the presence of 2  $\mu$ l of each of the dNTPs  
at 10 mM. The reaction was carried out at 37°C for 10 min. The  
PCR product so treated, corresponding to the fragment «Tnos -  
LB» (477 bp), was then isolated by 2% agarose gel  
20 electrophoresis in TBE buffer and purified with the aid of the  
«QIAquick Gel Extraction» kit. The DNA was recovered in 30  $\mu$ l  
of H<sub>2</sub>O.

A second PCR amplification was then carried out from  
7  $\mu$ l of each of the PCR products treated, corresponding to  
25 «end nptII» and «Tnos - LB», with the aid of 20 pmoles of each  
of the oligodeoxynucleotides, 5'  
TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5'  
TTCTTGACGAGTTCTTCTGAGCGGG 3' (SEQ.ID34), in the presence of  
200  $\mu$ M of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH 9.1, 18 mM  
30 (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of ELONGASE enzyme (GIBCO BRL  
Life Technologies) in a final reaction medium of 50  $\mu$ l. The  
PCR amplification reaction was carried out in the «GeneAmp PCR  
System 9700» thermocycler. After denaturation at 94°C for 3  
min., the DNA was subjected to 15 cycles each made up of the  
35 stages of denaturation at 94°C for 45 sec., hybridization at  
60°C for 45 sec. and elongation at 68°C for 1 min. Then, in

the last cycle, the elongation was continued at 68°C for 3 min. This reaction is repeated 3 times for the PCR products treated, corresponding to «end nptII» and «Tnos - LB». The PCR reaction medium was then subjected to 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragment (672 bp), corresponding to «end nptII - Tnos - LB», was recovered in 100 µl of H<sub>2</sub>O. Next, 95 µl of this DNA were hydrolysed by AvrII, purified with the aid of the «QIAquick PCR Purification» kit, digested by BstBI, purified with the aid of the «QIAquick PCR Purification» kit and recovered in 50 µl of H<sub>2</sub>O. The 631 bp AvrII-BstBI fragment carries the sequence «end nptII - Tnos - LB».

3.2. Synthesis of the fragment carrying «end of the nos promoter (Pnos) and start of the nptII gene» («end Pnos - st. nptII»)

The fragment (1023 bp) carrying «end Pnos - st. nptII» was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5' GGAATCGAAATCTCGTGATGGCAGG 3' (SEQ.ID39) and 5' ATTATTGCGCGTTCAAAAGTCGCC 3' (SEQ.ID40), and that the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 min. The amplification was repeated for 2 samples of pBin19 DNA.

3.3. Synthesis of the fragment «end Pnos - nptII - Tnos - LB»

The fragments, «end Pnos - st. nptII» and «end nptII - Tnos - LB», were assembled by splicing overlap extension in order to produce a fragment «end Pnos - nptII - Tnos - LB».

To do this, a PCR amplification was carried out from 5 µl of each of the PCR products treated, corresponding to «end Pnos - st. nptII» and «end nptII - Tnos - LB», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5' ATTATTGCGCGTTCAAAAGTCGCC 3' (SEQ.ID40), in the presence of

200  $\mu$ M of each of the dNTPs, 60 mM Tris- $\text{SO}_4$  pH 9.1, 18 mM  $(\text{NH}_4)_2 \text{SO}_4$ , 1.8 mM  $\text{MgSO}_4$  and 2 U of  $\text{ELONGASE}$  enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50  $\mu$ l. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 3 min. This reaction was repeated 5 times for each of the PCR products treated, corresponding to «end Pnos - st. nptII» and «end nptII - Tnos - LB». The PCR reaction medium was then subjected to 0.7% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragment (1608 bp) corresponding to «end Pnos - nptII - Tnos - LB», was recovered in 50  $\mu$ l of  $\text{H}_2\text{O}$ . Next, this DNA was subjected to the action of 62.5 U of the Klenow fragment (New England Biolabs) in the presence of 15  $\mu$ l of the Klenow fragment x 10 buffer (New England Biolabs) and 10  $\mu$ l of each of the dNTPs at 10 mM. The reaction was carried out at 37°C for 10 min. The DNA so treated was then purified with the aid of the «QIAquick PCR Purification» kit and recovered in 50  $\mu$ l of  $\text{H}_2\text{O}$ .

#### 3.4. Synthesis of the fragment carrying «RB - MCS - st. Pnos»

3.4.1. Obtaining the fragment carrying «RB and multiple cloning site (MCS)» («RB - MCS»)

The fragment (210 bp) carrying «RB - MCS» was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5' CGGTACCGAAGCTTTGAATTCACTCGAGCAGATTGTCGTTTCCCGCC 3' (SEQ.ID35) possessing the restriction sites KpnI, HindIII, EcoRI and XhoI, and 5' TATCCTAGGAACCGGTAAACCCTGTGGTTGGCATGC 3' (SEQ.ID36) possessing the restriction sites AvrII and AgeI.

3.4.2. Obtaining the fragment carrying «multiple cloning site (MCS) and start of the nos promoter (Pnos)» («MCS - st. Pnos»)

The fragment (209 bp) carrying «MCS - st. Pnos» was amplified and treated in the same way as the fragment carrying LB except that the 2 oligodeoxynucleotides used are 5' ATATGAGACTCTAATTGGATACCGAGGGG 3' (SEQ.ID37) and 5' GCTCGAGTGAATTCAAAGCTTCGGTACCGTTGAAGGAGCCACTCAGCCG 3' (SEQ.ID38) possessing the restriction sites XhoI, EcoRI, HindIII and KpnI.

3.4.3. Obtaining the fragment carrying «RB - MCS - st. Pnos»

The fragments, «RB - MCS» and «MCS - st. Pnos», were assembled by splicing overlap extension in order to produce a fragment «RB - MCS - st. Pnos».

To do this, a PCR amplification was carried out from 12 µl of each of the PCR products treated, corresponding to «RB - MCS» and «MCS - st. Pnos», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' ATATGAGACTCTAATTGGATACCGAGGGG 3' (SEQ.ID37) and 5' TATCCTAGGAACCGGTAAACCCTGTGGTTGGCATGC 3' (SEQ.ID36), in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 30 sec., hybridization at 60°C for 30 sec. and elongation at 68°C for 45 sec. Then, in the last cycle, the elongation was continued at 68°C for 3 min. This reaction was repeated for the remaining 3 x 12 µl of each of the PCR products treated, corresponding to «RB - MCS» and «MCS - st. Pnos». The PCR reaction medium was then subjected to 2% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit (QIAGEN). The DNA fragment

(390 bp) corresponding to «RB - MCS - st. Pnos», was recovered in 100 µl of H<sub>2</sub>O. Next, 95 µl of this DNA was hydrolysed by AvrII, purified with the aid of the «QIAquick PCR Purification» kit, digested by Bsu36I, purified with the aid of the «QIAquick PCR Purification» kit and recovered in 50 µl of H<sub>2</sub>O. The 295 bp AvrII-Bsu36I fragment carries the sequence «RB - MCS - st. Pnos».

### 10 3.5 Synthesis of the T-DNA

The fragments, «RB - MCS - st. Pnos» and «end Pnos - nptII - Tnos - LB», were assembled by splicing overlap extension in order to produce a fragment «RB - MCS - Pnos - nptII - Tnos - LB» which corresponds to the T-DNA (1883 bp).

15 To do this, a PCR amplification was carried out from 4 µl of the PCR product treated corresponding to «RB - MCS - st. Pnos» and from 5 µl of the PCR product treated corresponding to «end Pnos - nptII - Tnos - LB», with the aid of 20 pmoles of each of the oligodeoxynucleotides, 5' TTCCTAGGTTGACGTCTTCTGATGGGCTGCCTGTATCG 3' (SEQ.ID29) and 5' TATCCTAGGAACCGGTAAACCCTGTGGTTGGCATGC 3' (SEQ.ID36), in the presence of 200 µM of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH 9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of ELONGASE enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 25 50 µl. The PCR amplification reaction was carried out in the «GeneAmp PCR System 9700» thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 30 min. Then, in the last cycle, the elongation was continued at 68°C for 3 min. This reaction was repeated 10 times for each of the PCR products treated, corresponding to «RB - MCS - st. Pnos» and «end Pnos - nptII - Tnos - LB». The PCR reaction medium was then subjected to 0.8% agarose gel electrophoresis 35 in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragment (1883 bp) corresponding to



the T-DNA was recovered in 100  $\mu$ l of H<sub>2</sub>O. Next, this T-DNA was hydrolysed by AvrII (1873 bp fragment), purified with the aid of the «QIAquick PCR Purification» kit and recovered in 100  $\mu$ l of H<sub>2</sub>O.

5                    3.6. Obtaining pMRT1118

The binary plasmid pMRT1118 (5971 bp) results from the introduction of the T-DNA fragment digested by AvrII into the AvrII site of the dephosphorylated pMRT1106 plasmid.

To do this, the pMRT1106 plasmid DNA (5  $\mu$ g) was  
10 digested by AvrII, purified with the aid of the «QIAquick PCR Purification» kit, then dephosphorylated by 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, isolated  
15 by 0.6% agarose gel electrophoresis in TBE buffer, purified with the aid of the «QIAquick Gel Extraction» kit, dephosphorylated a second time with the calf intestinal alkaline phosphatase enzyme under the aforesaid conditions, and finally purified with the aid of the «QIAquick PCR  
20 Purification» kit and recovered in 50  $\mu$ l of H<sub>2</sub>O.

The ligation by PCR reaction was carried out with 32.5 ng of digested dephosphorylated pMRT1106 plasmid and 50 ng of digested T-DNA fragments in a reaction medium of 10  $\mu$ l in the presence of 1  $\mu$ l of T4 DNA ligase x 10 buffer (New  
25 England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec., in the «GeneAmp PCR System 9700» thermocycler.

The bacteria, *Escherichia coli* DH5 $\alpha$ , which had  
30 previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The  
35 resultant plasmid selected was called pMRT1118 (Figure 3). Its complete sequence SEQ.ID03 is given in the sequence listing.

#### 4. Synthesis of pMRT1119

The plasmid pMRT1119 (6016 bp) differs from pMRT1118 by the addition of additional unique restriction sites in the multiple cloning site MCS of pMRT1118.

##### 4.1. Obtaining the additional unique restriction sites

The additional unique restriction sites (XbaI, SalI, PacI, BamHI, MluI, HpaI and FseI) were created by hybridization between the 2 oligodeoxynucleotides, 5'AGCTTGGCCGGCCGTTAACAACGGTGGATCCTTAATTAAAGTCGACTCTAGAG 3' (SEQ.ID41) and 5' AATTCTCTAGAGTCGACTTAAATTAAAGGATCCACGGTGTAAACGGCCGCCA 3' (SEQ.ID42). To do this, 5 µg of each of the 2 oligodeoxynucleotides were mixed and held at 85°C for 1 min., followed by a progressive reduction of the temperature to 80°C for 5 min., then a slow reduction of the temperature to 60°C in a water bath, and finally a rapid reduction of the temperature to the ambient temperature outside the water bath.

##### 4.2. Obtaining pMRT1119

The binary plasmid pMRT1119 (6016 bp) results from the introduction of the sequence carrying the unique restriction sites into the HindIII and EcoRI sites of the pMRT1118 plasmid.

To do this, the pMRT1118 plasmid DNA (5 µg) was doubly digested by HindIII and EcoRI, purified with the aid of the «QIAquick PCR Purification» kit, and recovered in 50 µl of H<sub>2</sub>O.

The ligation by PCR reaction was carried out with 75 ng of digested pMRT1118 plasmid and 500 ng of fragments carrying the unique restriction sites (described in 4.1.) in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec. in the «GeneAmp PCR System 9700» thermocycler.

The bacteria, *Escherichia coli* DH5 $\alpha$ , which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and verified by enzymatic digestions and by sequencing. The resultant plasmid selected was called pMRT1119 (Figure 4). Its complete sequence SEQ.ID04 is given in the sequence listing.

10           5. Synthesis of pMRT1121

The plasmid pMRT1121 (6017 bp) results from the insertion into pMRT1119 of a unique restriction site, BspEI, between the nopaline synthase promoter of *Agrobacterium tumefaciens* (Pnos) and the mutant nptII gene.

15           The BspEI site was inserted by the assembly by splicing overlap extension of two fragments obtained by PCR amplification.

          5.1. Synthesis of the fragment carrying "part of nptII and BspEI site"

20           The 892 bp fragment carrying "part of nptII and BspEI site" was amplified by PCR from 5 ng of pBin19 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' GGAATCGAAATCTCGTGATGGCAGG 3' (SEQ.ID39) and 5' TAATCTGCATCCGGATCTGGATCGTTTCGC 3' (SEQ.ID43) carrying the BspEI site, in the presence of 200  $\mu$ M of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH 9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of eLONGase enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50  $\mu$ l. This reaction was repeated. The PCR amplification reaction was carried out in a "GeneAmp PCR 25 System 9700" thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 55°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 3 30 min. The PCR product so obtained was isolated by 2% agarose gel electrophoresis in TBE buffer and purified with the aid of

the «QIAquick Gel Extraction» kit. The DNA was recovered in 50  $\mu$ l of H<sub>2</sub>O.

## 5.2. Synthesis of the fragment carrying "BspEI site - MCS"

5 The fragment (250 bp) carrying "BspEI site - MCS" was amplified by PCR from 5 ng of pMRT1118 matrix DNA with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' GCTCGAGTGAATTCAAAGCTTCGGTACCGTTGAAGGAGCCACTCAGCCG 3' (SEQ.ID38) and 5' ACGATCCAGATCCGGATGCAGATTATTTGG 3' (SEQ.ID44) carrying the BspEI site. The conditions of PCR amplification and treatment of the PCR fragment obtained are the same as described in 5.1.

## 15 5.3. Synthesis of the fragment carrying "part nptII - MCS"

The 1117 bp fragment carrying "part nptII - MCS" results from the assembly of the 2 PCR fragments, "part nptII - BspEI site" and "BspEI site - MCS", by splicing overlap extension.

To do this, a PCR amplification was carried out from 7.5  $\mu$ l of each of the PCR products treated, corresponding to "part nptII - BspEI site" and "BspEI site - MCS", with the aid of 20 pmoles of each of the 2 oligodeoxynucleotides, 5' GGAATCGAAATCTCGTGATGGCAGG 3' (SEQ.ID39) and 5' GCTCGAGTGAATTCAAAGCTTCGGTACCGTTGAAGGAGCCACTCAGCCG 3' (SEQ.ID38), in the presence of 200  $\mu$ M of each of the dNTPs, 60 mM Tris-SO<sub>4</sub> pH 9.1, 18 mM (NH<sub>4</sub>)<sub>2</sub> SO<sub>4</sub>, 1.8 mM MgSO<sub>4</sub> and 2 U of eLONGase enzyme (GIBCO BRL Life Technologies) in a final reaction medium of 50  $\mu$ l. This reaction was repeated 4 times. The PCR amplification reaction was carried out in a "GeneAmp PCR System 9700" thermocycler. After denaturation at 94°C for 3 min., the DNA was subjected to 15 cycles each made up of the stages of denaturation at 94°C for 45 sec., hybridization at 62°C for 45 sec. and elongation at 68°C for 1 min. Then, in the last cycle, the elongation was continued at 68°C for 7

min. The PCR product so obtained was isolated by 1% agarose gel electrophoresis in TBE buffer and purified with the aid of the "QIAquick Gel Extraction" kit. The DNA was recovered in 50  $\mu$ l of H<sub>2</sub>O, digested by Bsu36I and PstI, and subjected to 2% agarose gel electrophoresis in TEB buffer. The 315 bp DNA fragment was isolated and purified with the aid of the "QIAquick Gel Extraction" kit.

#### 5.4. Obtaining pMRT1121

The 315 bp DNA fragment Bsu36I - PstI, which carries the BspEI site, was ligated to the Bsu36I and PstI sites of the binary plasmid pMRT1119.

The plasmid pMRT1119 was previously digested by Bsu36I and PstI, subjected to 1% agarose gel electrophoresis in TEB buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA was recovered in 50  $\mu$ l of H<sub>2</sub>O, dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the "QIAquick PCR Purification" kit.

The ligation by PCR reaction was carried out with 100 ng of digested dephosphorylated pMRT1119 plasmid and 30 ng of digested DNA fragments (315 bp) in a reaction medium of 10  $\mu$ l in the presence of 1  $\mu$ l of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec., in a "GeneAmp PCR System 9700" thermocycler. The bacteria, Escherichia coli DH5 $\alpha$  or SCS110 (deficient in Dam and Dcm methylases), which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1121 (Figure 5). Its complete sequence SEQ.ID05 is given in the sequence listing.

The plasmid pMRT1122 (6016 bp) results from the introduction into pMRT1119 of a point mutation in the mutant nptII gene in order to restore the BglII restriction site and thus lead to the obtaining of the wild-type nptII gene.

10 The oligodeoxynucleotide.  
5' ATGGGTACACGACGAGATCTTCGCCGTCGGG 3' (SEQ.ID45) was previously  
phosphorylated by subjecting 600 pmoles of the  
oligodeoxynucleotide to the action of 90 units of T4 kinase  
(Amersham) in the presence of 30  $\mu$ l of T4 kinase x 10 buffer  
15 (Amersham) and 3  $\mu$ l of 100 mM ATP in a final reaction medium  
of 300  $\mu$ l at 37°C for 30 min. The oligodeoxynucleotide so  
treated was purified with the aid of the «Qiaquick Removal  
Nucleotide» (QIAGEN) kit in accordance with the supplier's  
recommendations.

30

hybridization at 50°C for 1 min. and elongation at 65°C for 4 min. ; and, finally, the last phase consisted of a cycle made up of the stages of denaturation at 94°C for 30 sec., hybridization at 50°C for 1 min. and elongation at 65°C for 15 min. The PCR reaction medium was then subjected to 0.8% agarose gel electrophoresis in TBE buffer and purified with the aid of the «QIAquick Gel Extraction» kit. The DNA fragments (1023 bp) so treated were hydrolysed by NcoI and PstI and subjected to 2% agarose gel electrophoresis. The 383 bp DNA fragments were isolated, purified with the aid of the «QIAquick Gel Extraction» kit, recovered in 50 µl of H<sub>2</sub>O and ligated to the NcoI and PstI sites of the pMRT1119 plasmid. To do this, the pMRT1119 plasmid DNA was digested by NcoI and PstI, purified on 0.8% agarose gel. The fragment corresponding to the plasmid was isolated , purified with the aid of the «QIAquick Gel Extraction» kit, then dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and, finally, recovered in 50 µl of H<sub>2</sub>O.

The ligation by PCR was carried out with 50 ng of digested dephosphorylated pMRT1119 plasmid and all the DNA fragments digested and treated in a reaction medium of 10 µl in the presence of 1 µl of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec., in the «GeneAmp PCR System 9700» thermocycler.

The bacteria, *Escherichia coli* DH5α, which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and screened by digestion by BglII. The plasmid DNA of the clone selected was verified by enzymatic digestions and by sequencing. The resultant plasmid selected was called pMRT1122

(6016 bp). It is represented in Figure 6 and its complete sequence SEQ.ID06 is given in the sequence listing.

#### 7. Synthesis of pMRT1155

5 The plasmid pMRT1155 (6017 bp) differs from pMRT1122 by the presence of the BspEI site.

To do this, the 315 bp insert fragment containing the BspEI site was obtained by digestion of pMRT1121 by Bsu36I and PstI, 1.2% agarose gel electrophoresis in TEB buffer, 10 purification with the aid of the "QIAquick Gel Extraction" kit and recovery in 50  $\mu$ l of H<sub>2</sub>O.

The pMRT1122 vector fragment was obtained by digestion of pMRT1122 by Bsu36I and PstI, 1.2% agarose gel electrophoresis in TEB buffer, purification with the aid of 15 the "QIAquick Gel Extraction" kit and recovery in 50  $\mu$ l of H<sub>2</sub>O. After that, digested pMRT1122 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 20 hour, and purified with the aid of the "QIAquick PCR Purification" kit.

The ligation by PCR reaction was carried out with 100 ng of dephosphorylated digested pMRT1122 plasmid and 50 ng of digested DNA fragments (315 bp) in a reaction medium of 10  $\mu$ l 25 in the presence of 1  $\mu$ l of T4 DNA ligase x 10 buffer (New England Biolabs) and 400 units of T4 DNA ligase enzyme (New England Biolabs). It is made up of 180 cycles consisting of 2 stages, one of 10°C for 30 sec. and the other of 30°C for 30 sec. in a "GeneAmp PCR System 9700" thermocycler. The 30 bacteria, Escherichia coli DH5 $\alpha$  or SCS110 (deficient in Dam and Dcm methylases), which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis 35 method and verified by enzymatic digestions and sequencing.



The resultant plasmid was called pMRT1155 (Figure 7). Its complete sequence SEQ.ID07 is given in the sequence listing.

8. Obtaining basic binary plasmids comprising the  
5 sequence "eP35S - T35S"

#### 8.1. Obtaining pMRT1205

The plasmid pMRT1205 (7503 bp) includes the expression cassette of the mutant nptII gene and a "double 35S promoter - 35S terminator" sequence (eP35S - T35S) for the  
10 cloning of genes of interest. It results from the cloning of the eP35S promoter of the cauliflower mosaic virus (CaMV) into pMRT1175. The CaMV eP35S promoter corresponds to a duplication of the transcription-activating sequences situated upstream of the TATA element of the 35S promoter (Kay et al., 1987). As  
15 for the plasmid pMRT1175 (6767 bp), this results from the cloning of T35S of the cauliflower mosaic virus into pMRT1121. The transcription-terminating sequence, CaMV T35S, corresponds to the non-coding region in 3' of the circular double-stranded DNA sequence of the cauliflower mosaic virus producing the 35S  
20 transcript (Franck et al., 1980).

#### 8.1.1. Obtaining pMRT1175

The plasmid pMRT1175 results from the cloning of the EcoRI - XhoI insert DNA fragments corresponding to the T35S  
25 into the EcoRI and XhoI sites of the vector pMRT1121 produced from Escherichia coli strain SCS110.

The pMRT1121 vector fragment was obtained by digestion of 7 µg of pMRT1121 by EcoRI and XhoI, purification with the aid of the "QIAquick PCR Purification" kit and recovery in 50 µl of H<sub>2</sub>O. After that, digested pMRT1121 was  
30 dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl  
35 of H<sub>2</sub>O.

The EcoRI - XhoI insert DNA fragments (757 bp) corresponding to the T35S (750 bp) were obtained by digestion by EcoRI and XhoI of the plasmid pJIT163, which derives from the plasmid pJIT60 (Guerineau and Mullineaux, 1993). They were  
5 then subjected to 1% agarose gel electrophoresis in TEB buffer, purified with the aid of the «QIAquick Gel Extraction» kit and recovered in 30 µl of H<sub>2</sub>O.

The ligation by PCR reaction was carried out as described earlier in 7, with 80 ng of dephosphorylated  
10 digested pMRT1121 plasmid and 80 ng of digested insert DNA fragments. The bacteria, Escherichia coli DH5α, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis  
15 method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1175 (6767 bp). It is represented in Figure 8 and its complete sequence SEQ.ID08 is given in the sequence listing.

20

#### 8.1.2. Obtaining pMRT1205

The plasmid pMRT1205 results from the cloning of the KpnI - HindIII insert DNA fragments corresponding to eP35S into the KpnI and HindIII sites of the vector pMRT1175.

25 The pMRT1175 vector fragment was obtained by digestion of 4.1 µg of pMRT1175 by KpnI and HindIII, purification with the aid of the «Concert Rapid PCR Purification System» kit and recovery in 100 µl of H<sub>2</sub>O. After that, digested pMRT1175 was dephosphorylated with 50 units of  
30 calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120 µl in the presence of 12 µl of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50 µl of H<sub>2</sub>O.

35 The KpnI - HindIII insert DNA fragments (743 bp) corresponding to eP35S (735 bp) were obtained by digestion by

KpnI and HindIII of the plasmid pJIT163, which derives from the plasmid pJIT60 (Guerineau and Mullineaux, 1993). They were then subjected to 1% agarose gel electrophoresis in TEB buffer, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 30  $\mu$ l of H<sub>2</sub>O.

The ligation by PCR reaction was carried out as described earlier in 7, with 48 ng of dephosphorylated digested pMRT1175 plasmid and 30 ng of digested insert DNA fragments. The bacteria, *Escherichia coli* DH5 $\alpha$ , which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1205 (7503 bp). It is represented in Figure 19 and its complete sequence SEQ.ID19 is given in the sequence listing.

#### 8.2. Obtaining pMRT1203

The plasmid pMRT1203 (7503 bp) includes the expression cassette of the wild-type nptII gene and a "double 35S promoter - 35S terminator" sequence (eP35S - T35S) for the cloning of genes of interest. It results from the cloning of the eP35S promoter of the cauliflower mosaic virus into pMRT1176. As for the plasmid pMRT1176 (6767 bp), this results from the cloning of T35S of the cauliflower mosaic virus into pMRT1155. The eP35S promoter and the T35S terminator are described in 8.1.

##### 8.2.1. Obtaining pMRT1176

The plasmid pMRT1176 results from the cloning of the EcoRI - XhoI insert DNA fragments (757 bp) corresponding to the T35S (750 bp) into the EcoRI and XhoI sites of the vector pMRT1155 produced from *Escherichia coli* strain SCS110.

The plasmid pMRT1176 was obtained in accordance with the methodologies described in 8.1.1., except that 6.3  $\mu$ g of the vector pMRT1155, which constitutes the cloning vector, were digested and treated. The plasmid pMRT1176 (6767 bp) is

represented in Figure 9. Its complete sequence SEQ.ID09 is given in the sequence listing.

#### 8.2.2. Obtaining pMRT1203

The plasmid pMRT1203 results from the cloning of the  
5 KpnI - HindIII insert DNA fragments (743 bp) corresponding to eP35S (735 bp) into the KpnI and HindIII sites of the vector pMRT1176.

The plasmid pMRT1203 was obtained in accordance with the methodologies described in 8.1.2., except that 3.9 µg of  
10 the vector pMRT1176, which constitutes the cloning vector, were digested and treated. The plasmid pMRT1203 (7503 bp) is represented in Figure 17. Its complete sequence SEQ.ID17 is given in the sequence listing.

#### 15 9. Obtaining binary plasmids comprising the sequence "eP35S - uidA - T35S"

These vectors were employed in transformation of tobacco in order to evaluate them and determine the incidence of the mutation of the nptII gene.

#### 20 9.1. Obtaining pMRT1206

The plasmid pMRT1206 (9390 bp) includes the expression cassette of the mutant nptII gene and the expression cassette of the uidA gene (gus). It results from the cloning of the eP35S promoter of the cauliflower mosaic  
25 virus into pMRT1196. As for the plasmid pMRT1196 (8654 bp), this results from the cloning of the uidA gene (Jefferson RA et al., 1986) into pMRT1175.

#### 9.1.1. Obtaining pMRT1196

The plasmid pMRT1196 results from the cloning of the  
30 insert DNA fragments (SmaI - "SacI + T4 DNA polymerase"), corresponding to the uidA gene, into the "XbaI + Klenow" site of the vector pMRT1175.

The pMRT1175 vector fragment was obtained by digestion of 10 µg of pMRT1175 by XbaI, purified with the aid  
35 of the "QIAquick PCR Purification" kit, recovered in 50 µl of H2O and subjected to the action of 20 units of the Klenow

fragment (New England Biolabs) in the presence of 12  $\mu$ l of 500 mM Tris-HCl pH 7.5, 500 mM  $MgCl_2$ , 6  $\mu$ l of 1M dithiothreitol, 6  $\mu$ l of each of the 10 mM dNTPs in a reaction volume of 120  $\mu$ l at 37°C for 30 min. It was then purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50  $\mu$ l of  $H_2O$ . The plasmid pMRT1175 so digested and treated was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, and purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50  $\mu$ l of  $H_2O$ .

The insert DNA fragments (2  $\mu$ g) corresponding to the uidA gene (1.8 kbp) were obtained by digestion of pBI221 (marketed by Clontech) by SacI, purification with the aid of the "QIAquick PCR Purification" kit and recovery in 50  $\mu$ l of  $H_2O$ . After that, digested pBI221 was subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of T4 DNA polymerase x 10 buffer, 4  $\mu$ l of 10 mM dNTPs and 6  $\mu$ l of BSA 1 mg/ml. The reaction was carried out at 37°C for 30 min. The plasmid pBI221 so treated was purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50  $\mu$ l of  $H_2O$ . Finally, pBI221 so treated was digested by SmaI. The «[SacI + T4 DNA polymerase] - SmaI» fragment (1882 bp) was isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 50  $\mu$ l of  $H_2O$ .

The ligation by PCR reaction was carried out in accordance with the methodologies described in 7, with 15 ng of dephosphorylated digested pMRT1175 plasmid and 100 ng of digested insert DNA fragments. The bacteria, Escherichia coli DH5 $\alpha$ , which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was

called pMRT1196 (8654 bp). It is represented in Figure 14 and its complete sequence SEQ.ID14 is given in the sequence listing.

#### 9.1.2. Obtaining pMRT1206

5. The plasmid pMRT1206 (9390 bp) results from the cloning of the KpnI - HindIII insert DNA fragments (743 bp), corresponding to eP35S (735 bp), into the KpnI and HindIII sites of the vector pMRT1196.

10 The plasmid pMRT1206 was obtained in accordance with the methodologies described in 8.1.2., except that 2 µg of the vector pMRT1196, which constitutes the cloning vector, were digested and treated. It is represented in Figure 24 and its complete sequence SEQ.ID24 is given in the sequence listing.

15 The plasmid pMRT1206 was introduced into Agrobacterium tumefaciens strain LBA4404 by direct transformation in accordance with the procedure of Holsters et al. (1978).

#### 9.2. Obtaining pMRT1204

20 The plasmid pMRT1204 (9390 bp) includes the expression cassette of the wild-type nptII gene and the expression cassette of the uidA gene. It results from the cloning of the eP35S promoter of the cauliflower mosaic virus into pMRT1192. As for the plasmid pMRT1192 (8654 bp), this results from the cloning of the uidA gene into pMRT1176.

25

#### 9.2.1. Obtaining pMRT1192

30 The plasmid pMRT1192 results from the cloning of the insert DNA fragments (SmaI - "SacI + T4 DNA polymerase"), corresponding to the uidA gene, into the "XbaI + Klenow" site of the vector pMRT1176.

35 The plasmid pMRT1192 (8654 bp) was obtained in accordance with the methodologies described in 9.1.2., except that the cloning vector is pMRT1176. It is represented in Figure 11 and its sequence SEQ.ID11 is given in the sequence listing.

#### 9.2.2. Obtaining pMRT1204

The plasmid pMRT1204 (9390 bp) results from the cloning of the KpnI - HindIII insert DNA fragments (743 bp), corresponding to eP35S (735 bp), into the KpnI and HindIII sites of the vector pMRT1192.

The plasmid pMRT1204 was obtained in accordance with the methodologies described in 8.1.2., except that 2 µg of the vector pMRT1192, which constitutes the cloning vector, were digested and treated. It is represented in Figure 18 and its complete sequence SEQ.ID18 is given in the sequence listing.

The plasmid pMRT1204 was introduced into *Agrobacterium tumefaciens* strain LBA4404 by direct transformation in accordance with the procedure of Holsters et al. (1978).

#### 10. Obtaining binary plasmids containing the bar gene

The final plasmids were employed in transformation of maize and/or tobacco in order to evaluate them.

##### 10.1. Creation of pMRT1210

The plasmid pMRT1210 (10003 bp) includes the expression cassette of the bar gene and the expression cassette of the uidA gene. It results from the cloning of the expression cassette of the uidA gene into pMRT1195. As for the plasmid pMRT1195 (6865 bp), this results from the cloning of the sequence "promoter followed by intron 1 of the rice actin gene" (McElroy D et al., 1991) into pMRT1191. The plasmid pMRT1191 (4805 bp) derives from pMRT1119.

##### 10.1.1. Obtaining pMRT1191

The plasmid pMRT1191 (4805 bp) differs from pMRT1119 by the deletion of the nos promoter and mutant nptII gene sequences.

It was obtained by digestion by Bsp120I of 10 µg of pMRT1119, followed by purification with the aid of the "QIAquick PCR Purification" kit and recovery in 50 µl of H<sub>2</sub>O. After that, digested pMRT1119 and [sic] was hydrolysed by

KpnI, purified with the aid of the "QIAquick PCR Purification" kit, recovered in 50  $\mu$ l of H<sub>2</sub>O, subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of T4 DNA polymerase x 10 buffer, 4  $\mu$ l of 10 mM dNTPs and 6  $\mu$ l of BSA at 1 mg/ml. The reaction was carried out at 37°C for 30 min. The vector fragment was then isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 50  $\mu$ l of H<sub>2</sub>O.

10           Thirty nanograms of the plasmid so treated were religated by ligation by PCR reaction (procedure described in 7).

          The resultant plasmid was called pMRT1191. It is represented in Figure 10 and its complete sequence SEQ.ID10 is  
15   given in the sequence listing.

#### 10.1.2. Obtaining pMRT1201

          The plasmid pMRT1201 (7943 bp) differs from pMRT1191 by the insertion of the expression cassette of the uidA gene isolated from the plasmid pUC19-Phmwg-IA-uidA-Tnos which  
20   comprises the uidA expression cassette inserted into the EcoRI and HindIII sites of pUC19 (marketed by New England Biolabs). This plasmid results from successive clonings. The fragment corresponding to the sequence «uidA gene - nos terminator» was isolated by successive digestions by «EcoRI followed by the  
25   action of the Klenow fragment» and SmaI from pBI221  $\Delta$  SacI (= pBI221 (Clontech) whose ScaI site was eliminated by SacI digestion followed by the action of T4 DNA polymerase, then ligation). The isolated and purified fragment was inserted into the «SmaI - [SphI + T4 DNA polymerase]» sites of pUC19  
30   (Clontech) in order to obtain the resultant plasmid pUC19-uidA-Tnos. After that, the EcoRI site recreated on the 3' side of Tnos was eliminated by replacement of the BstBI-HindIII fragment of pUC19-uidA-Tnos with the BstBI-HindIII fragment obtained by digestion of the fragment amplified by PCR (1054  
35   bp) from the pUC19-uidA-Tnos matrix with the aid of the 2 oligodeoxynucleotides 5' AGGCATTGGTTTCGAAGCG 3' (SEQ.ID46)



containing the BstBI site and 5' TACGCCAAGCTTGGCAATTCC  
3' (SEQ.ID47). The resultant plasmid was called pUC19-uidA-Tnos  
ΔEcoRI. Then, in order to introduce a NcoI site at the level  
of the initiator codon of the uidA gene, a 440 bp fragment was  
5 amplified by PCR from the pUC19-uidA-TnosΔEcoRI matrix with  
the aid of the 2 oligodeoxynucleotides 5'  
AATACCCGGGACCATGGTCCGTCCTGTAG 3' (SEQ.ID48) containing the  
NcoI and SmaI sites and 5' ATAGTCTGCCAGTTCAGTTCGTTG 3'  
(SEQ.ID49) situated downstream of SnaBI. The PCR fragment  
10 digested by SmaI and SnaBI was then inserted into pUC19-uidA-  
Tnos Δ EcoRI in order to replace the existing SmaI-SnaBI  
fragment. The resultant plasmid was called pUC19-NcoI-uidA-  
Tnos. After that, the Phmwg promoter [High Molecular Weight  
Glutenin, Anderson et al. (1989)], carried by the EcoRI - SmaI  
15 fragment, was introduced into the EcoRI and SmaI sites of  
pUC19-NcoI-uidA-Tnos in order to produce the plasmid pUC19-  
Phmwg-uidA-Tnos. Finally, the intron 1 of the rice actin gene  
(IA, EcoRV - SmaI fragment) comes from the plasmid pAct1-F6  
(McElroy D et al., 1991) which is itself modified by the  
20 addition of restriction sites notably on both sides of the  
EcoRV-SmaI fragment containing IA. The SmaI - NcoI fragment  
carrying IA was isolated from pACT1-F6 modified and inserted  
into the SmaI and NcoI sites of pUC19-Phmwg-uidA-Tnos in order  
to produce the plasmid pUC19-Phmwg-IA-uidA-Tnos. The  
25 techniques used are the ones already described in this patent.

The pMRT1191 vector fragment was obtained by  
digestion of 2 μg of pMRT1191 by EcoRI, purification with the  
aid of the «Concert Rapid PCR Purification System» kit,  
followed by the action of 20 units of the Klenow fragment (New  
30 England Biolabs) in the presence of 12 μl of 500 mM Tris-HCl  
pH 7.5, 500 mM MgCl<sub>2</sub>, 6 μl of 1M dithiothreitol, 6 μl of each  
of the dNTPs at 10 mM in a reaction volume of 120 μl at 37°C  
for 30 min., purification with the aid of the «Concert Rapid  
PCR Purification System» kit, and recovery in 50 μl of H<sub>2</sub>O.  
35 After that, digested and treated pMRT1191 was dephosphorylated

with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid  
5 PCR Purification System» kit, and recovered in 50  $\mu$ l of H<sub>2</sub>O.

The insert DNA fragments (2  $\mu$ g) corresponding to the expression cassette of the uidA gene (3 kbp), viz. the uidA gene placed under the control of the wheat "high molecular weight glutenin" promoter followed by the intron 1 of the rice  
10 actin gene and the nopaline synthase terminator of Agrobacterium tumefaciens, were obtained by digestion of pUC19-Phmwg-IA-uidA-Tnos by ScaI (site present in pUC19), purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50  $\mu$ l of H<sub>2</sub>O, digestion by HindIII,  
15 purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50  $\mu$ l of H<sub>2</sub>O, treatment by 20 units of the Klenow fragment (New England Biolabs) in the presence of 12  $\mu$ l of 500 mM Tris-HCl, pH 7.5, 500 mM MgCl<sub>2</sub>, 6  $\mu$ l of 1M dithiothreitol, 6  $\mu$ l of each of the 10 mM dNTPs in a reaction  
20 volume of 120  $\mu$ l at 37°C for 30 min., purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50  $\mu$ l of H<sub>2</sub>O and digestion by EcoRI. After that, the insert DNA fragments were isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit,  
25 subjected to the action of the Klenow fragment as described above, and recovered in 50  $\mu$ l of H<sub>2</sub>O.

The ligation by PCR reaction was carried out as described earlier in 7, with 10 ng of dephosphorylated digested pMRT1191 plasmid and 100 ng of digested and treated  
30 insert DNA fragments. The bacteria, Escherichia coli DH5 $\alpha$ , which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and  
35 sequencing. The resultant plasmid was called pMRT1201 (7943

bp). It is represented in Figure 15 and its complete sequence SEQ.ID15 is given in the sequence listing.

#### 10.1.3. Obtaining pMRT1210

The plasmid pMRT1210 (10003 bp) differs from pMRT1201  
5 by the insertion of the sequence "promoter followed by the  
intron 1 of the rice actin gene - bar gene" (Pact-IA-bar)  
isolated from pSB12-Pact-IA-bar-Tnos which derives from pSB12  
described by Komari et al. (1996). The plasmid pSB12-Pact-IA-  
bar-Tnos results from the cloning of the expression cassette  
10 «Pact-IA-bar-Tnos» (BspDI - «XhoI + Klenow» fragment),  
isolated from pDM302, into the SmaI and BspDI sites of pSB12,  
whose XhoI site on the LB side was deleted. The plasmid  
pDM302, constructed in the laboratory of Wu R., comprises the  
expression cassette «Pact-IA-bar-Tnos» in the plasmid pSP72  
15 marketed by Promega.

The pMRT1201 vector fragment was obtained by  
digestion of 2 µg of pMRT1201 by HpaI, purification with the  
aid of the «Concert Rapid PCR Purification System» kit, and  
recovery in 98 µl of H<sub>2</sub>O. After that, digested pMRT1201 was  
20 dephosphorylated with 50 units of calf intestinal alkaline  
phosphatase (New England Biolabs) in a final reaction medium  
of 120 µl in the presence of 12 µl of 3 x 10 buffer (New  
England Biolabs) at 37°C for 1 hour, purified with the aid of  
the «Concert Rapid PCR Purification System» kit, and recovered  
25 in 50 µl of H<sub>2</sub>O.

The insert DNA fragments (2 µg) corresponding to  
"Pact-IA-bar" (2.1 kbp) were obtained by XbaI digestion of  
pBIOS273, purification with the aid of the "QIAquick PCR  
Purification" kit, recovery in 50 µl of H<sub>2</sub>O, treatment with 20  
30 units of the Klenow fragment (New England Biolabs) in the  
presence of 12 µl of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl<sub>2</sub>,  
6 µl of 1M dithiothreitol and 6 µl of each of the 10 mM dNTPs  
in a reaction volume of 120 µl at 37°C for 30 min. The insert  
DNA fragments so treated were then isolated by 0.8% agarose  
35 gel electrophoresis, purified with the aid of the "QIAquick  
PCR Purification" kit and recovered in 50 µl of H<sub>2</sub>O.

The ligation by PCR reaction was carried out as described earlier in 7, with 60 ng of dephosphorylated digested pMRT1201 plasmid and 100 ng of digested and treated insert DNA fragments. The bacteria, *Escherichia coli* DH5 $\alpha$ , which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1210. It is represented in Figure 21 and its complete sequence SEQ.ID21 is given in the sequence listing.

This plasmid pMRT1210 was then introduced into *Agrobacterium tumefaciens* strain LBA4404 (pSB1) [Komari T et al., 1996] and strain LBA4404 by direct transformation in accordance with the procedure of Holsters et al. (1978).

#### 10.1.4. Obtaining pMRT1193

The binary plasmid pMRT1193 was obtained by cloning, into the EcoRI and [XhoI + Klenow fragment action] sites of pMRT1119, of the «EcoRI - [HindIII + Klenow fragment action]» fragment carrying the sequence «Phmwg-IA-uidA-Tnos» isolated from pUC19-Phmwg-IA-uidA-Tnos, described in 10.1.2.

To do this, the pMRT1119 vector fragment was obtained by digestion of 10  $\mu$ g of pMRT1119 by XhoI, purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50  $\mu$ l of H<sub>2</sub>O, treatment with 20 units of the Klenow fragment (New England Biolabs) in the presence of 12  $\mu$ l of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl<sub>2</sub>, 6  $\mu$ l of 1M dithiothreitol and 6  $\mu$ l of each of the 10 mM dNTPs in a reaction volume of 120  $\mu$ l at 37°C for 30 min. Then purification with the aid of the "QIAquick PCR Purification" kit and digestion by EcoRI. The treated vector fragment was isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50  $\mu$ l of H<sub>2</sub>O. After that, digested pMRT1119 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3

x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the "QIAquick PCR Purification" kit and recovered in 50 µl of H<sub>2</sub>O.

The insert DNA fragments (2 µg) corresponding to the expression cassette of the uidA gene (3 kbp), viz. the uidA gene placed under the control of the wheat "high molecular weight glutenin" promoter followed by the intron 1 of the rice actin gene and the nopaline synthase terminator of *Agrobacterium tumefaciens*, were obtained by digestion of pUC19-Phmwg-IA-uidA-Tnos by ScaI (site present in pUC19), purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H<sub>2</sub>O, digestion by HindIII, purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H<sub>2</sub>O, treatment with 20 units of the Klenow fragment (New England Biolabs) in the presence of 12 µl of 500 mM Tris-HCl, pH 7.5, 500 mM MgCl<sub>2</sub>, 6 µl of 1M dithiothreitol and 6 µl of each of the 10 mM dNTPs in a reaction volume of 120 µl at 37°C for 30 min., purification with the aid of the "QIAquick PCR Purification" kit, recovery in 50 µl of H<sub>2</sub>O, and digestion by EcoRI. The insert DNA fragments were then isolated by 0.8% agarose gel electrophoresis, purified with the aid of the "QIAquick Gel Extraction" kit and recovered in 50 µl of H<sub>2</sub>O.

The ligation by PCR reaction was carried out as described earlier in 7, with 20 ng of dephosphorylated digested pMRT1119 plasmid and 100 ng of digested and treated insert DNA fragments. The bacteria, *Escherichia coli* DH5α, which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1193 (9143 bp). It is represented in Figure 12 and its complete sequence SEQ.ID12 is given in the sequence listing.

35

#### 10.2. Obtaining pMRT1195

The plasmid pMRT1195 (6857 bp) differs from pMRT1191 by the insertion of the sequence "promoter followed by the intron 1 of the rice actin gene - bar gene" (Pact-IA-bar). The vector pMRT1191 was digested by Bsp120I and KpnI followed by  
5 the action of the enzyme T4 DNA polymerase and dephosphorylated but not religated was obtained as described in 10.1.1.

The insert DNA fragments corresponding to "Pact-IA-bar" (2.1 kbp) were obtained as described in 10.1.3.

10 The ligation by PCR reaction was carried out as described earlier in 7, with 20 ng of dephosphorylated digested pMRT1191 plasmid and 100 ng of digested and treated insert DNA fragments. The bacteria, *Escherichia coli* DH5a, which had previously been made competent, were transformed.  
15 The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1195. It is represented in Figure 13 and its complete sequence SEQ.ID13 is  
20 given in the sequence listing.

#### 10.3. Obtaining pMRT1202

The plasmid pMRT1202 (5614 bp) differs from pMRT1195 by the replacement of the sequence "Pact-IA" by the nopaline synthase (Pnos) promoter of *Agrobacterium tumefaciens* isolated  
25 from pMRT1121 produced in *Escherichia coli* strain SCS110.

The pMRT1195 vector fragment (2  $\mu$ g) was digested by PstI, purified with the aid of the «Concert Rapid PCR Purification System» kit, recovered in 98  $\mu$ l of H<sub>2</sub>O, treated by the action of 6 units of T4 DNA polymerase (New England  
30 Biolabs) in a reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of T4 DNA polymerase x 10 buffer, 4  $\mu$ l of 10 mM dNTPs and 6  $\mu$ l of BSA at 1 mg/ml. The reaction was carried out at 37°C for 30 min. The vector fragment was then purified with the aid of the «Concert Rapid PCR Purification System» kit,  
35 recovered in 50  $\mu$ l of H<sub>2</sub>O, digested by HindIII, isolated by 0.8% agarose gel electrophoresis, purified with the aid of the

«Concert Rapid PCR Purification System» kit, and recovered in 98  $\mu$ l of H<sub>2</sub>O. After that, digested and treated pMRT1195 was dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50  $\mu$ l of H<sub>2</sub>O.

The insert DNA fragments corresponding to "Pnos" were obtained by BspEI digestion of pMRT1121, purification with the aid of the «Concert Rapid PCR Purification System» kit, recovery in 98  $\mu$ l of H<sub>2</sub>O, treatment with 20 units of the Klenow fragment (New England Biolabs) in the presence of 12  $\mu$ l of 500 mM Tris-HCl, pH 7.5 - 500 mM MgCl<sub>2</sub>, 6  $\mu$ l of 1M dithiothreitol and 6  $\mu$ l of each of the 10 mM dNTPs in a reaction volume of 120  $\mu$ l at 37°C for 30 min., purification with the aid of the «Concert Rapid PCR Purification System» kit, recovery in 50  $\mu$ l of H<sub>2</sub>O, and digestion by HindIII. The insert DNA fragments (219 bp) carrying Pnos (209 bp) were then isolated by 2% agarose gel electrophoresis, purified with the aid of the «Concert Rapid PCR Purification System» kit, and recovered in 50  $\mu$ l of H<sub>2</sub>O.

The ligation by PCR reaction was carried out as described earlier in 7, with 48 ng of dephosphorylated digested pMRT1195 plasmid and 56 ng of digested and treated insert DNA fragments. The bacteria, Escherichia coli DH5 $\alpha$ , which had previously been made competent, were transformed. The plasmid DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1202. It is represented in Figure 16 and its complete sequence SEQ.ID16 is given in the sequence listing.

#### 10.4. Obtaining pMRT1212

The plasmid pMRT1212 (8987 bp) differs from pMRT1206 by the replacement of the nptII gene by the bar gene.

To do this, the pMRT1206 vector fragment was obtained by digestion by Bsu36I of 2  $\mu$ g of pMRT1206, purification with the aid of the «Concert Rapid PCR Purification System» kit, recovery in 50  $\mu$ l of H<sub>2</sub>O and digestion by AatII. The digestion  
5 by Bsu36I and AatII allowed deletion of the vector of the fragment corresponding to «part Pnos nptII Tnos LB». The digested vector fragment was then subjected to 1% agarose gel electrophoresis, purified with the aid of the «Concert Rapid PCR Purification System» kit and recovered in 98  $\mu$ l of H<sub>2</sub>O.  
10 This digested and treated vector fragment was then dephosphorylated with 50 units of calf intestinal alkaline phosphatase (New England Biolabs) in a final reaction medium of 120  $\mu$ l in the presence of 12  $\mu$ l of 3 x 10 buffer (New England Biolabs) at 37°C for 1 hour, purified with the aid of  
15 the «Concert Rapid PCR Purification System» kit and recovered in 50  $\mu$ l of H<sub>2</sub>O.

The insert DNA fragments corresponding to «part Pnos bar Tnos LB» (1.2 kbp) were obtained by digestion by Bsu36I of 2  $\mu$ g of pMRT1202, purification with the aid of the Concert  
20 Rapid PCR Purification System kit, recovery in 40  $\mu$ l of H<sub>2</sub>O, digestion by AatII, isolation by 1% agarose gel electrophoresis, purification with the aid of the Concert Rapid PCR Purification System kit and recovery in 50  $\mu$ l of H<sub>2</sub>O.

25 The ligation by PCR reaction was carried out as described earlier in 7, with 100 ng of dephosphorylated digested pMRT1206 plasmid and 50 ng of digested insert DNA fragments. The bacteria, Escherichia coli DH5 $\alpha$ , which had previously been made competent, were transformed. The plasmid  
30 DNA of the clones obtained, selected on LB medium supplemented with kanamycin (50 mg/l), was extracted by the alkaline lysis method and verified by enzymatic digestions and sequencing. The resultant plasmid was called pMRT1212. It is represented in Figure 22 and its complete sequence SEQ.ID22 is given in  
35 the sequence listing.



11. Evaluation of the synthetic vectors produced in the bacteria

The yield of plasmids obtained from the bacteria *Escherichia coli* strain DH5 $\alpha$  was determined.

5 I Non-binary synthetic plasmids

Primary cultures of the recombinant bacteria possessing the plasmids pMRT1105, pMRT1106 or pMRT1105-ori ColE1 were prepared from 500 ml of each of the respective "glycerol-stocks" in 10 ml of LB medium supplemented with  
10 kanamycin at 50 mg/l, at 37°C for 16 hours. After that, 500 ml of each of the primary cultures was cultured in 100 ml of LB supplemented with kanamycin at 50 mg/l at 37°C for 23 hours. For pMRT1106, two cultures were prepared, one without addition of chloramphenicol and the other with addition of  
15 chloramphenicol at 40 mg/l after 7 hours of culture. The cultures were then centrifuged at 5000 g for 10 min. The plasmid DNAs were extracted with the aid of the "QIAfilter Plasmid Midi kit" (QIAGEN), in accordance with the manufacturer's recommendations, and quantitatively determined  
20 with the spectrophotometer. The quantities of plasmids obtained were determined as 10 mg, 50.6 mg, 86.2 mg and 9.8 mg respectively for pMRT1105, pMRT1106, pMRT1106 treated with chloramphenicol, and pMRT1105 ori ColE1 (plasmid pMRT1105 in which ori ColE1 was inserted in the reverse orientation to  
25 that of pMRT1106). These results show that the synthetic plasmids are replicated in *Escherichia coli*, the presence of the origin of replication "ori ColE1" in the right orientation (NdeI site of "ori ColE1" near "ori RK2") allows the yield to be increased by a factor of 5, whereas no effect is shown for  
30 the reverse orientation of "ori ColE1", treatment with chloramphenicol contributes an additional factor of 1.6.

11.2. Synthetic binary plasmids.

The synthetic binary plasmids pMRT1118 and pMRT1119 are evaluated relative to the controls pBin19 and pBIOC4. The  
35 plasmid pBIOC4 differs from pGA492 (An, 1986) by the deletion of virtually all the coding sequence of the pGA492 cat gene

and by the conversion of the HindIII site into an EcoRI site. It was obtained by double digestion by SacI and ScaI followed by the action of the enzyme T4 DNA polymerase (New England Biolabs) and ligation of the modified plasmid (20 ng) in a reaction medium of 10  $\mu$ l in the presence of 1  $\mu$ l of T4 DNA ligase x 10 buffer (Amersham) and 2.5 units of T4 DNA ligase (Amersham) at 14°C for 16 hours. The bacteria, *Escherichia coli* strain DH5 $\alpha$ , which had previously been made competent, were transformed (Hanahan, 1983). The plasmid DNA of the clones obtained, selected on tetracycline at 12 mg/l, was extracted by the alkaline lysis method (Birnboim and Doly, 1979) and analysed by enzymatic digestions. The HindIII site of the plasmid DNA of the clone selected were then modified into an EcoRI site with the aid of a phosphorylated HindIII-EcoRI adaptor (Stratagene). To do this, 500 ng of plasmid DNA of the clone selected were digested by HindIII, dephosphorylated by the enzyme calf intestinal alkaline phosphatase (Boehringer Mannheim) in accordance with the manufacturer's recommendations and coprecipitated in the presence of 1500 ng of HindIII-EcoRI adaptor DNA, 0.1 volume of 3M sodium acetate, pH 4.8, and 2.5 volumes of absolute ethanol at -80°C for 30 min. After centrifugation at 12000 g for 30 min., the precipitated DNA was washed with 70% ethanol, dried, recovered in 8  $\mu$ l of H<sub>2</sub>O, held at 65°C for 10 min., then ligated as described earlier. After inactivation of the T4 DNA ligase at 65°C for 10 min., the ligation reaction mixture was digested by EcoRI, purified by 0.8% agarose gel electrophoresis, electroeluted, precipitated with absolute ethanol, centrifuged at 12000 g for 30 min., washed with 70% ethanol, dried, then ligated and introduced into *Escherichia coli* strain DH5 $\alpha$ , and treated as described earlier.

Primary cultures of the recombinant bacteria possessing the plasmids mentioned above were prepared from 500 ml of each of the respective "glycerol-stocks" in 10 ml of LB medium supplemented with kanamycin at 50 mg/l, except for pBIOC4, which was selected on tetracycline at 12 mg/l, at 37°C.

for 16 hours. After that, 250 ml of each of the primary cultures were cultured in 100 ml of LB supplemented with the appropriate antibiotic at 37°C for 23 hours. The cultures were then centrifuged at 5000 g for 10 min. The plasmid DNAs were  
5 extracted with the aid of the "QIAfilter Plasmid Midi kit" (QIAGEN) in accordance with the manufacturer's recommendations and quantitatively determined with the spectrophotometer. The quantities of plasmids obtained were determined as 94.2 mg, 113 mg, 37.2 mg and 45.6 mg respectively for pMRT1118,  
10 pMRT1119, pBin19 and pBIOC4.

These results show that the synthetic binary plasmids are replicated in *Escherichia coli*, and that the yield of pMRT1119 is increased by the factors 3 and 2.5 respectively, relative to the pBin19 and pBIOC4 controls.

15 Furthermore, these synthetic plasmids are replicated in *Agrobacterium tumefaciens* strain LBA4404.

12. Evaluation of the synthetic plasmids by transgenesis

12.1. Transformation of tobacco

20 12.1.1. Stable transformation of tobacco

The transformation of tobacco (*Nicotiana tabacum* L. var. PBD6) was carried out by infecting foliar discs isolated from 6-week-old tobacco seedlings in vitro with recombinant agrobacteria in accordance with the method described by Horsch  
25 et al. (1985).

All the in vitro cultures are prepared in a controlled environment chamber in which the light intensity is 200  $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ , the photoperiod 16 hours light/8 hours darkness, and the temperature 25°C.

30 Except for the initial coculture stage, the regeneration, development and rooting stages were performed on different selective media supplemented with a bacteriostat, namely augmentin at 400 mg/l, and a selective agent, namely kanamycin at 200 or 100 mg/l.

35 The different stages and the media used are as follows:

After primary culture of the agrobacteria in 5 ml of 2YT medium (bacto-tryptone 10 g/l, yeast extract 5 g/l, NaCl 5 g/l, pH 7.2) supplemented with  $\text{CaCl}_2$ , with a final concentration of 6 mM, and with appropriate antibiotics, at 28°C for 48 hours, a culture is performed in 10 ml of 2YT medium, supplemented with  $\text{CaCl}_2$  and antibiotics, at 28°C for a period of one night. The culture is then centrifuged at 3000 g for 10 min. and the bacteria are resuspended in 10 ml of liquid MS30 (M0404 at 4.4 g/l, marketed by SIGMA, supplemented with sucrose, 30 g/l, pH 5.7).

The coculture is performed by placing the foliar explants cut from the leaves of the seedlings in vitro, about 1 cm<sup>2</sup>, in contact with the suspension of agrobacteria, diluted 1/10, in liquid MS30 for 20 min. The explants so treated are then rapidly dried on filter paper and placed on a solid coculture medium (CM) (MS30, Benzyl Amino Purine (BAP) 1 mg/l, Indole-3 Acetic Acid (ANA) at 0.1 mg/l, agar at 8 g/l) for 48 hours in the controlled environment chamber.

The treated explants are then placed on a solid regeneration medium (solid CM, augmentin 400 at mg/l, kanamycin at 200 mg/l). The explants are pricked out on the same medium after 2 weeks.

After 2 weeks the shoots are pricked out on a solid development medium (M0404 at 4.4 g/l, marketed by SIGMA, supplemented with sucrose 20 g/l, pH 5.7 (liquid MS20), augmentin at 400 mg/l, kanamycin at 100 mg/l, agar at 8 g/l).

After 2 weeks, the transformed seedlings are pricked out on a solid rooting medium identical to the development medium. Rooting takes 2 to 3 weeks, after which the seedlings are removed to the growth chamber in Giffy pots for 10 days (photoperiod 16 hours light/8 hours darkness, 23°C, 70% relative humidity), then placed in the greenhouse.

12.1.2. Evaluation of the synthetic plasmids used in transformation of tobacco.

The recombinant agrobacteria (*Agrobacterium tumefaciens* strain LBA4404) containing the synthetic binary

plasmid pMRT1118 or the control binary plasmid pBin19 were used to transform the tobacco *Nicotiana tabacum* L. var. PBD6.

The results showed that the number and growth of the seedlings which develop in the presence of the selective agent (kanamycin) are similar for the two binary plasmids tested. These observations indicate that the synthetic binary plasmid is perfectly functional in transgenesis of tobacco.

## 12.2. Transformation of maize

### 12.2.1. Genetic transformation of maize calli

The genetic transformation of maize, whatever method is employed (electroporation, *Agrobacterium*, microfibres, particle gun), generally requires the use of cells that are undifferentiated in rapid divisions, having retained the ability to regenerate whole plants. Cells of this type make up the friable embryogenic callus (called Type II) of maize.

These calli were obtained from immature embryos of genotype HI, II or (A188 x B73) in accordance with the method and on the media described by Armstrong (Maize handbook, 1994, Freeling M and Walbot V (eds), pp 665-671). The calli so obtained were multiplied and maintained by successive subcultures every 2 weeks on the starting medium.

Seedlings were regenerated from these calli by modifying the hormonal and osmotic equilibrium of the cells in accordance with the method described by Vain et al. (1989). These plants were then hardened off in the greenhouse, where they can be crossed or selfed.

### 12.2.2. Genetic transformation of maize with the particle gun

The production and regeneration of the cell lines needed for transformation are described in 12.2.1.

The procedure which uses the particle gun for genetic transformation is described by Finer et al. (1992). The target cells are callus fragments with a surface area of 10 to 20 mm<sup>2</sup>. These fragments were placed on a starting medium supplemented with 0.2 M mannitol and 0.2 M sorbitol for 4 hours before bombardment.

The tungsten particles (M10) and transformation plasmids were coprecipitated in accordance with the protocol described by Klein (1987). The particles thus coated were projected towards the target cells by means of the particle gun in accordance with the protocol of Finer et al. (1992).

The bombarded calli were placed in darkness at 27°C. After 24 hours, the first subculturing took place, followed by subculturing every 2 weeks for 3 months on a starting medium to which the appropriate selective agent to select only the transformed calli had been added. These calli were then grown in the presence of the selective agent in order to regenerate transformed seedlings, as described in 12.2.1. The seedlings obtained were hardened off and transferred to the greenhouse, where they can be crossed or selfed.

#### 12.2.3. Genetic transformation of maize by *Agrobacterium tumefaciens*

The technique used is described by Ishida et al. (1996). Immature embryos 1.0 to 1.2 mm in length (9 to 14 days after pollination) were washed in the LS-inf medium, then immersed in the suspension of agrobacteria, prepared as described by Ishida et al. (1996), vortexed for 30 sec., and incubated at ambient temperature for 5 min. The immature embryos so treated were cultured on LS-AS medium in darkness at 25°C for 3 days, then transferred to LSD 1.5 medium supplemented with phosphinothricin at 5 mg/l and cefotaxime at 250 mg/l, in darkness at 25°C for 2 weeks, and finally placed on LSD 1.5 medium supplemented with phosphinothricin at 10 mg/l and cefotaxime at 250 mg/l, in darkness at 25°C for 3 weeks. The Type I calli thus generated were isolated, fragmented and transferred to LSD 1.5 medium supplemented with phosphinothricin at 10 mg/l and cefotaxime at 250 mg/l, in darkness at 25°C for 3 weeks. The Type I calli which proliferated were then isolated and placed on LSZ medium supplemented with phosphinothricin at 5 mg/l and cefotaxime at 250 mg/l, subjected to a photoperiod of 16 hours light/8 hours darkness at 25°C for 2 to 3 weeks. The regenerated seedlings

were then transferred to ¼ LSF medium subjected to a photoperiod of 16 hours light/8 hours darkness at 25°C for 1 to 2 weeks, and then transferred to the growth chamber and greenhouse.

5                   13. Influence of the vector backbone on gene expression

Stably transformed tobacco plants were obtained for the synthetic binary plasmids pMRT1204 bearing the wild-type nptII gene (16 plants) and pMRT1206 bearing the mutated nptII gene  
10 (12 plants), and also for the control vector pBI121 sold by Clontech which corresponds to plasmid pBIN19 bearing the wild-type nptII gene and containing the expression cassette "P35S - uidA - polyA nos" (11 plants).

The influence of the synthetic vector backbone and that of the  
15 control was evaluated by ELISA analysis of protein extracts using the NPTII ELISA kit sold by 5 Prime -> 3 Prime. The analysis enabled a quantification of the NPTII protein produced (in micrograms, µg) with respect to the amount of soluble extracted protein (in milligrams, mg).

20 The proteins were extracted in "Tris-HCl 25 mM pH7.8 ; Phenylmethylsulfonylfluoride 1 mM" buffer from tobacco leaves ground in liquid nitrogen. After centrifugation for 10 minutes at 10000 g and 4°C, the extracted soluble proteins contained in the supernatant were measured according to Bradford's  
25 method (A rapid and sensitive method for the detection of microgram quantities of proteins utilizing the principle of protein-dye binding. Anal. Biochem. (1976) 72, 248-254).

The antibodies used for ELISA quantification were rabbit polyclonal antibodies specifically recognizing the coating  
30 NPTII protein and biotinylated anti-NPTII antibodies for the detection of the NPTII protein.

The results of the analysis shown in Figure 23 indicate that :

for the wild type nptII gene cassette and different plasmid  
35 backbones, the quantities of NPTII obtained with the synthetic binary plasmid pMRT1204 are greater by a factor

of 6.7 than those obtained with the original pBIN19 binary plasmid, which demonstrates the efficiency of the synthetic vector.

5 for an identical plasmid backbone, i.e. synthetic binary plasmids pMRT1204 and pMRT1206, but with different nptII genes, i.e. Wild type and mutated nptII, the quantities of NPTII obtained in the wild type gene differ from those obtained for the mutated nptII gene by a factor 1.2.

#### 14. Synthesis of pMRT1334.

10 The plasmid pMRT1334 (9688 bp) was obtained by replacing the nptII expression cassette of pMRT1206 by the nptII expression cassette of pBIN19.

The vector fragment derived from pMRT1206 was obtained by digestion of 10  $\mu$ g of pMRT1206 with KpnI, purified  
15 using a "Concert Rapid PCR Purification System" kit, and taken up in 50  $\mu$ l of water. Then the digested pMRT1206 was subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction mixture of 120  $\mu$ l in the presence of 12  $\mu$ l of T4 10x DNA polymérase buffer, 4  $\mu$ l of 10 mM dNTP and  
20 6  $\mu$ l of 1 mg/ml BSA. The reaction was carried out at 37°C for 30 minutes. The digested and thus treated pMRT1206 vector was purified with a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l of water, then digested with AflIII. The vector fragment derived from pMRT1206 was isolated by  
25 electrophoresis on a 1% agarose gel, purified with a "Concert Rapid PCR Purification System" and taken up in 50  $\mu$ l of water.

The DNA insert fragment DraI - AflIII (1.5 kbp), corresponding to the nptII expression cassette was obtained by digestion of 9  $\mu$ g of pBIN19 plasmid with DraI and AflIII. Then,  
30 the fragment was subjected to electrophoresis on a 1% agarose gel in TEB buffer, purified using a "Concert Rapid PCR Purification System", and taken up in 50  $\mu$ l of water.

The PCR ligation reaction was carried out using 100 ng of the plasmid fragment derived from pMRT1206 and 100  
35 ng of the DNA insert fragment DraI - AflIII in a reaction mixture of 20  $\mu$ l in the presence of 2  $\mu$ l of T4 10x DNA ligase



buffer (Epicentre Technologies), 2  $\mu$ l 2.5 mM ATP and 4 units of T4 DNA ligase (Epicentre Technologies). The reaction consists of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a  
5 "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent *Escherichia coli* DH5 $\alpha$  bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline  
10 lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1334 (Figure 24). Its complete sequence is given under SEQ.ID50 in the sequence listing.

#### 15. Synthesis of pMRT1335

15 The plasmid pMRT1335 (15208 bp) is a control vector and results from the insertion of the expression cassette "ep35S-gus (uidA)-polyA35S" isolated from pMRT1206 into pBIN19.

13  $\mu$ g of pBIN19 plasmid were digested with EcoRI,  
20 purified using a "Concert Rapid PCR Purification System", and taken up in 50  $\mu$ l water. Then the previously digested pBIN19 was subjected to the action of 20 units of Klenow fragment (New England Biolabs) in the presence of 12  $\mu$ l 500 mM pH7.5 Tris-HCl 500 mM MgCl<sub>2</sub>, 6  $\mu$ l of 1M dithiothreitol, 6  $\mu$ l each of  
25 10 mM dNTP in a reaction volume of 120  $\mu$ l at 37°C for 60 minutes. The digested pBIN19 vector was purified using a "Concert Rapid PCR Purification System", taken up in 50  $\mu$ l water, then digested with KpnI. The vector fragment thus  
30 produced was isolated by electrophoresis on a 1% agarose gel, and purified using a "Concert Rapid PCR Purification System" and taken up in 50  $\mu$ l of water.

The DNA insert fragment corresponding to the "ep35S-gus-polyA35S" (3.5 kbp) expression cassette was obtained by digestion of 10  $\mu$ g of plasmid pMRT1206 with XhoI, purified  
35 using a "Concert Rapid PCR Purification System", and taken up in 50  $\mu$ l water, followed by the action of 20 units of Klenow

fragment (New England Biolabs) in the presence of 12  $\mu$ l of 500 mM pH7.5 Tris-HCl 500 mM MgCl<sub>2</sub>, 6  $\mu$ l 1M dithiothreitol, 6  $\mu$ l each of 10 mM dNTP in a reaction volume of 120  $\mu$ l at 37°C for 60 minutes, then purified using a "Concert Rapid PCR Purification System", taken up in 50  $\mu$ l water, and finally digested with KpnI. The DNA insert fragment thus produced was isolated by electrophoresis on a 1% agaros gel in TEB buffer, and purified using a "Concert Rapid PCR Purification System", and taken up in 50  $\mu$ l water.

10 The PCR ligation reaction was carried out with 100 ng of pBIN19 treated and 100 ng of DNA insert fragment as prepared previously in a reaction volume of 20  $\mu$ l in the presence of 2  $\mu$ l of T4 10x DNA ligase buffer (Epicentre Technologies), 2  $\mu$ l 2.5 mM ATP and 4 units T4 DNA ligase 15 (Epicentre Technologies). The reaction consists of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent *Escherichia coli* DH5 $\alpha$  20 bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1335 25 (Figure 25). Its complete sequence is given under SEQ.ID51 in the sequence listing.

#### 16. Synthesis of pMRT1336

The plasmid pMRT1336 (9285 bp) results from the insertion into pMRT1196 of the promoter MPr1165 (610 bp) 30 isolated from plasmid pMRT1322 as described in PCT patent application PCT/IB00/00370, and incorporated into the present description by reference thereto for that relevant part.

10  $\mu$ g of pMRT1196 plasmid were digested with KpnI, purified using a "Concert Rapid PCR Purification System", taken up in 50  $\mu$ l water, and redigested with HpaI. The thus 35 treated vector fragment was subjected to electrophoresis on a

1% agarose gel, and purified using a "Concert Rapid PCR Purification System" and then taken up in 50  $\mu$ l water.

The DNA insert fragment bearing the promoter MPrl165 (0.5 kbp) was obtained from 10  $\mu$ g of pMRT1322 plasmid digested with KpnI, purified using a "Concert Rapid PCR Purification System" and taken up in 50  $\mu$ l water, and then redigested with HpaI. The DNA insert fragment thus produced was subjected to gel electrophoresis on a 1% agarose gel in TEB buffer, purified using a "Concert Rapid PCR Purification System", and taken up in 50  $\mu$ l water.

The PCR ligation reaction was carried out using 100 ng of vector fragment and 100 ng DNA insert fragment bearing the promoter MPrl165 as prepared above in a reaction volume of 20  $\mu$ l in the presence of de 2  $\mu$ l T4 10x DNA ligase buffer (Epicentre Technologies), 2  $\mu$ l 2.5 mM ATP and 4 units T4 DNA ligase (Epicentre Technologies). The reaction consists of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent Escherichia coli DH5 $\alpha$  bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1336 (Figure 26). Its complete sequence is given in the sequence listing under SEQ.ID52.

Plasmid pMRT1322 was obtained by the insertion of promoter MPrl165 into pMRT1176. 10  $\mu$ g of pMRT1176 plasmid was digested with XbaI and EcoRI, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, dephosphorylated with 50 units of calf intestine alkaline phosphatase (New England Biolabs) in a final reaction volume of 120  $\mu$ l in the presence 12  $\mu$ l of 3x10 buffer (New England Biolabs) at 37 °C for 1 hour, and purified using a "Concert Rapid PCR Purification System" kit, and then taken up in 50  $\mu$ l

water. The DNA insert fragment corresponding to promoter MPr1165 was obtained by PCR amplification from 10 ng of pMRT1240 matrix DNA using 20 pmoles each of two oligodeoxynucleotides, 5' AGCTCTAGAGCTGCCTGCAGCACTAGTATCC  
5 3' (SEQ.ID53) bearing the site XbaI and 5' CGGAATTCGGCCTCTAGGTTGTTGTGTTG 3' (SEQ.ID54) bearing the site EcoRI, using the enzyme Platinum Taq polymerase High Fidelity (GIBCO BRL Life Technologies) and following the suppliers instructions. The PCR amplification reaction was carried out  
10 in a "GeneAmp PCR System 9700" thermocycler. After denaturation at 94°C for 2 minutes, the DNA was subjected to 25 cycles, each including denaturation steps at 94°C for 45 seconds, hybridisation at 55°C for 45 seconds and elongation at 68°C for 45 seconds. On the final cycle the elongation was  
15 continued at 68°C for 3 minutes. The PCR product obtained was isolated by electrophoresis on a 1.5% agarose gel in TBE buffer and purified using a "Concert Rapid PCR Purification System" kit, and then taken up in 50 µl water. The PCR product was then digested with XbaI and EcoRI, purified using a  
20 "Concert Rapid PCR Purification System" kit, and taken up in 50 µl water. The PCR ligation reaction was carried out as described previously with 80 ng of pMRT1176 vector fragment and 110 ng of DNA insert fragment bearing the MPr1165 promoter. Previously prepared competent Escherichia coli DH5α  
25 bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1322.  
30 Plasmid pMRT1240 was obtained by insertion of a DNA fragment into plasmid pMRT1234, by treatment with Klenow fragment (New England Biolabs) according to the supplier's instructions of the HindIII sites and BamHI. The DNA insert fragment was the L5 promoter fragment obtained by double PstI  
35 digestion, and BamHI digestion, followed by the action of T4 DNA polymerase (New England Biolabs) according to the

supplier's instructions, of plasmid pMRT1165, as described in PCT patent application PCT/IB00/00370, and incorporated into the present description by reference thereto for that relevant part.

5 Plasmid pMRT1234 was obtained by the insertion at the EcoRI site of pMRT1175 of a DNA insert fragment EcoRI corresponding to the cDNA coding for human lactoferrin isolated from the plasmid pHMWG-IA-PSSp-Lf, as described in previously published patent application WO98/50543.

10 17. Synthesis of pMRT1337

Plasmid pMRT1337 (8289 bp) was obtained from the insertion, into pMRT1205, of the gfp gene isolated from pBINm-gfp5-ER and described in Haseloff J et Siemering KR. 1998, "The uses of green fluorescent protein in plants", in Green fluorescent protein : Strategies, applications and protocols (Chalfie M et Kain S, eds, Wiley, pp 191-220).

15 10  $\mu$ g of plasmid pMRT1205 was digested with EcoRI, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, subjected to the action of 20 units of Klenow fragment (New England Biolabs) in the presence of 20 12  $\mu$ l of 500 mM pH7.5 Tris-HCl 500 mM MgCl<sub>2</sub>, 6  $\mu$ l 1M dithiothreitol, 6  $\mu$ l each of 10 mM dNTP in a reaction volume of 120  $\mu$ l at 37°C for 60 minutes, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, 25 and then digested with BamHI. The thus treated vector fragment was subjected to electrophoresis on 1% agarose gel, purified using a "Concert Rapid PCR Purification System" kit and taken up in 50  $\mu$ l water.

The DNA insert fragment bearing the gfp gene (0.7 30 kbp) was obtained from 10  $\mu$ g of plasmid pBINm-gfp5-ER digested with SacI, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, and subjected to the action of 6 units of T4 DNA polymerase (New England Biolabs) in a reaction volume of 120  $\mu$ l in the presence of 12  $\mu$ l T4 10x 35 DNA polymerase buffer, 4  $\mu$ l 10 mM dNTP and 6  $\mu$ l 1 mg/ml BSA. The reaction was carried out at 37°C for 30 minutes. The thus

treated plasmid pBINm-gfp5-ER was then purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, and digested with BamHI. The DNA insert fragment thus produced was subjected to gel electrophoresis on 1% gel agarose in TEB buffer, purified using a "Concert Rapid PCR Purification System" kit and taken up in 50  $\mu$ l in water.

The PCR ligation reaction was carried out with 100 ng of pMRT1205 vector fragment and 100 ng DNA insert fragment bearing the gfp gene as prepared above in a reaction volume of 20  $\mu$ l in the presence of 2  $\mu$ l T4 10x DNA ligase buffer (Epicentre Technologies), 2  $\mu$ l 2.5 mM ATP and 4 units T4 DNA ligase (Epicentre Technologies). The reaction comprised 180 cycles each including two steps, the first at 10°C for 30 seconds, the second at 30°C for 30 seconds, in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent Escherichia coli DH5 $\alpha$  bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1337 (Figure 27). Its complete sequence is given in the sequence listing under SEQ.ID55.

18. Synthesis of pMRT1341  
The plasmid pMRT1341 (14108 bp) results from the replacement of the "ep35S-gus-polyA35S" expression cassette from pMRT1335 by the expression cassette "ep35S-gfp-polyA35S" isolated from pMRT1337.

10  $\mu$ g of plasmid pMRT1335 was digested by AgeI and KpnI. The thus digested vector fragment was isolated by electrophoresis on a 0.8% agarose gel, purified using a "Concert Rapid PCR Purification System" kit and then take up in 50  $\mu$ l water.

The DNA insert fragment corresponding to the expression cassette "ep35S-gfp-polyA35S" (2.4 kbp) was obtained by digesting with XhoI of 10  $\mu$ g plasmid pMRT1337,

purifying using a "Concert Rapid PCR Purification System" kit, taking up in 50  $\mu$ l water, and the action of 20 units of Klenow fragment (New England Biolabs) in the presence of 12  $\mu$ l 500 mM pH7.5 Tris-HCl 500 mM MgCl<sub>2</sub>, 6  $\mu$ l 1M dithiothreitol, 5 6  $\mu$ l each of 10 mM dNTP in a reaction volume of 120  $\mu$ l at 37°C for 30 minutes, then, purifying using a "Concert Rapid PCR Purification System" kit, taking up in 50  $\mu$ l water, and finally digesting with KpnI. The thus prepared DNA insert fragment was isolated by electrophoresis on 0.8% agarose gel in TEB buffer, 10 purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, digested with AgeI, purified using a "Concert Rapid PCR Purification System" and taken up again in 50  $\mu$ l water.

The PCR ligation reaction was carried out on 100 ng 15 vector fragment derived from pMRT1335 et 100 ng DNA insert fragment as produced above in a reaction volume of 20  $\mu$ l in the presence of 2  $\mu$ l T4 10x DNA ligase buffer (Epicentre Technologies), 2  $\mu$ l 2,5 mM ATP and 4 units of T4 DNA ligase (Epicentre Technologies). The reaction comprised 180 cycles 20 each including 2 steps, the first at 10°C for 30 seconds, the second at 30°C for 30 seconds, in a "GeneAmp PCR System 9700" thermocycler.

Previously prepared competent *Escherichia coli* DH5 $\alpha$  bacteria, were transformed (Hanahan, 1983). The plasmid DNA of 25 the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and sequencing. The resulting plasmid was designated pMRT1341 (Figure 28). Its complete sequence is given in the sequence 30 listing under SEQ.ID56.

#### 19. Synthèse de pMRT1342

Plasmid pMRT1342 (15077 bp) results from the replacement of the expression cassette "ep35S-gus-polyA35S" from pMRT1335 by the expression cassette "L5-gus-polyA35S" 35 isolated from pMRT1336.

10  $\mu$ g of plasmid pMRT1335 was digested with AgeI and

KpnI. The thus digested vector fragment was isolated by electrophoresis on 0.8% agarose gel, purified using a "Concert Rapid PCR Purification System" kit and taken up in 50  $\mu$ l water.

5           The DNA insert fragment corresponding to the expression cassette "L5-gus-polyA35S" (3.4 kbp) was obtained by digestion with XhoI of 10  $\mu$ g plasmid pMRT1336, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, subjecting this to the action of 20 units  
10 Klenow fragment (New England Biolabs) in the presence of 12  $\mu$ l 500 mM pH7.5 Tris-HCl 500 mM MgCl<sub>2</sub>, 6  $\mu$ l 1M dithiothreitol, 6  $\mu$ l each of 10 mM dNTP in a reaction volume of 120  $\mu$ l at 37°C for 30 minutes, then, purified using a "Concert Rapid PCR Purification System" kit, and taken up in 50  $\mu$ l water, and  
15 finally digested with KpnI. The thus prepared DNA insert fragment was isolated by electrophoresis on a 0.8% agarose gel in TEB buffer, purified using a "Concert Rapid PCR Purification System" kit, taken up in 50  $\mu$ l water, digested with AgeI, purified using a kit as above and taken up again in  
20 50  $\mu$ l water.

The PCR ligation reaction was carried out with 100 ng of vector fragment derived from pMRT1335 and 100 ng of DNA insert fragment prepared as described above in reaction mixture of 20  $\mu$ l in the presence of 2  $\mu$ l T4 10x DNA ligase  
25 buffer (Epicentre Technologies), 2  $\mu$ l 2.5 mM ATP and 4 units T4 DNA ligase (Epicentre Technologies). The reaction is comprised of 180 cycles each including 2 steps, the first at 10°C for 30 seconds and the second at 30°C for 30 seconds in a "GeneAmp PCR System 9700" thermocycler.

30           Previously prepared competent Escherichia coli DH5 $\alpha$  bacteria, were transformed (Hanahan, 1983). The plasmid DNA of the obtained clones, selected on LB media supplemented with kanamycin (50 mg/l), was extracted according to the alkaline lysis method and verified by enzymatic digestion and  
35 sequencing. The resulting plasmid was designated pMRT1342 (Figure 29). Its complete sequence is given in the sequence



listing under SEQ.ID57.

20. Production of recombinant agrobacteria

5       The binary plasmids were transferred into the  
Agrobacterium tumefaciens LBA4404 strain according to the  
technique described by Holsters et al. [Holsters M., Dewaele  
D., Depicker A., Messenf E., Van Montagu M. et Schell J.  
(1978).. Transfection and transformation of Agrobacterium  
tumefaciens. Mol. Gen. Genet. 136, 181-187.]. The plasmid DNA  
10 of the obtained clones, selected on LB media supplemented with  
rifampicine (50 mg/l), was extracted according to the alkaline  
lysis method, modified by the addition of lysozyme (25 mg/ml)  
in the cell resuspension buffer. The plasmid DNA obtained was  
analysed by enzymatic digestion. The agrobacteria clones  
obtained were used for plant genetic transformation.

## References

- An G. 1986. Development of plant promoter expression  
5 vector and their use for analysis of differential activity of  
nopaline synthase promoter in transformed tobacco cells. *Plant  
Physiol.* 81 :86-91.
- Anderson OD, Green FC, Yip RE, Halford NG, Shewry PR  
and Malpica-Romero JM. 1989. Nucleotide sequences of the two  
10 high-molecular-weight glutenin genes from the D-genome of a  
hexaploid bread wheat, *Triticum aestivum* L. Cheyenne. *Nucl.  
Acids Res.* 17 :461-462.
- Birnboim HC and Doly J. 1979. A rapid alkaline  
extraction procedure for screening recombinant plasmid DNA.  
15 *Nucl. Acids Res.* 7 :1513.
- Depicker A, Stachel S, Dhaese P, Zambryski P and  
Goodman HM. 1982. Nopaline synthase : transcript mapping and  
DNA sequence. *J. Mol. Appl. Genet.* 1 :561-573.
- Finer JJ, Vain P, Jones MW and McMullen MD. 1992.  
20 Development of the particle inflow gun for DNA delivery to  
plant cells. *Plant Cell Rep.* 11 :323-328.
- Franck A, Guilley H, Jonard G, Richards K and Hirth  
L. 1980. Nucleotide sequence of cauliflower mosaic virus DNA.  
*Cell* 21 :285-294.
- 25 Frisch DA, Harris-Haller LW, Yokubaitis NT, Thomas  
TL, Hardin SH and Hall TC. 1995. Complete sequence of the  
binary vector Bin19. *Plant Mol. Biol.* 27 : 405-409.
- Guerineau F and Mullineaux P. 1993. Plant  
transformation and expression vectors. In *Plant Molecular*  
30 *Biology Labfax*. Croy RRD (ed). BIOS Scientific Publishers,  
Blackwell Scientific Publications, pp 121-147.
- Hanahan D. 1983. Studies on transformation of  
*Escherichia coli* with plasmids. *J. Mol. Biol.* 166 :557.
- Holsters M, Dewaele D, Depicker A, Messenf E, Van  
35 Montagu M and Schell J. 1978. Transfection and transformation  
of *Agrobacterium tumefaciens*. *Mol. Gen. Genet.* 136 :181-187.

- Horsch RB, Fry JE, Hoffmann NL, Eiholtz D, Rogers SG and Fraley RT. 1985. A simple and general method for transferring genes into plants. *Science* 227 :1229-1231.
- Ishida Y, Saito H, Ohta S, Hiei Y, Komari T and Kumashiro T. 1996. High efficiency transformation of maize (*Zea mays* L.) mediated by *Agrobacterium tumefaciens*. *Nature Biotechnology* 14 :745-750.
- Jefferson RA, Burgess SM and Hirsh D. 1986. b-glucuronidase from *Escherichia coli* as a gene-fusion marker. *Proc. Natl. Acad. Sci. (USA)* 83 :8447-8451.
- Kay R, Chan A, Daly M and McPherson J. 1987. Duplication of CaMV 35S promoter sequences creates a strong enhancer for plant genes. *Science* 236 :1299-1302.
- Klein TM, Wolf ED, Wu R and Sanford JC. 1987. High velocity microprojectiles for delivering nucleic acids into living cells. *Nature* 327 :70-73.
- Komari T, Hiei Y, Yasuhito S, Murai N and Kumashiro T. 1996. Vectors carrying two separate T-DNAs for co-transformation of higher plants mediated by *Agrobacterium tumefaciens* and segregation of transformants free from selection markers. *Plant J.* 10 :165-174.
- McElroy D, Blowers AD, Jenes B and Wu R. 1991. Construction of expression vectors based on the rice actin 1 (Act1) 5' region for the use in monocot transformation. *Mol. Gen. Genet.* 231 :150-160.
- Michael SF. 1994. Mutagenesis by incorporation of a phosphorylated oligo for PCR amplification. *BioTechniques* 16 :411-412.
- Vain P, Yean H and Flament P. 1989. Enhancement of production and regeneration of embryogenic type II callus in *Zea mays* L. by  $\text{AgNO}_3$ . *Plant Cell Tissue and Organ Culture* 18 :143-151.